

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 December 2001 (06.12.2001)

PCT

(10) International Publication Number
WO 01/92330 A2

- (51) International Patent Classification⁷: **C07K 14/47**
- (21) International Application Number: **PCT/IB01/01126**
- (22) International Filing Date: **22 May 2001 (22.05.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
60/208,251 31 May 2000 (31.05.2000) US
60/239,735 11 October 2000 (11.10.2000) US
- (71) Applicant (for all designated States except US): **GENSET** [FR/FR]; Intellectual Property Department, 24, rue Royale, F-75008 Paris (FR).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **LODISH, Harvey** [US/US]; 195 Fisher Avenue, Brookline, MA 02146 (US). **FRUEBIS, Joachim** [DE/US]; 2433 Summerhill Drive, Encinitas, CA 92024 (US). **TSAO, Tsu-Shuen** [US/US]; 453 Broadway #3, Somerville, MA 02145 (US). **BIHAIN, Bernard** [BE/US]; 238 Pacific View Lane, Encinitas, CA 92024 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **USE OF ACRP30 GLOBULAR HEAD TO PROMOTE INCREASES IN MUSCLE MASS AND MUSCLE DIFFERENTIATION**

(57) Abstract: The present invention relates to the field of muscle research, in particular to the discovery of a compound effective for increasing muscle mass, muscle cell differentiation, and oxidation of free fatty acids in muscle, useful in methods of treating muscle-related diseases and disorders as well as for augmenting muscle mass in general. The muscle-related diseases or disorders envisaged to be treated by the methods of the invention include, but are not limited to, muscular dystrophy, and other conditions resulting in muscle atrophy or muscle wasting.

WO 01/92330 A2

USE OF ACRP30 GLOBULAR HEAD TO PROMOTE INCREASES IN MUSCLE MASS AND MUSCLE DIFFERENTIATION

FIELD OF THE INVENTION

5 The present invention relates to the field of muscle research, in particular to the discovery of a compound effective for increasing muscle mass, muscle cell differentiation, and oxidation of free fatty acids in muscle, that should be useful in methods of treating muscle-related diseases and disorders as well as for augmenting muscle mass. The muscle-related diseases or disorders envisaged to be treated by the methods of the invention include, but are not limited to, muscle-related eye diseases and
10 disorders, muscle-related recovery after injuries, muscle-related recovery after surgery, muscle-related disorders of aging, muscular dystrophy, and other conditions resulting in muscle atrophy.

BACKGROUND OF THE INVENTION

15 Acrp30 (also known as AdipoQ) is the murine homolog of the human Apm1 protein (this class of proteins is generically referred to as OBG3 herein). The predicted protein structure of Apm1 indicates the presence of: 1) a secreted protein signal peptide, 2) a region containing collagen repeats, and 3) a globular region (Fig. 2), which is highly conserved (Fig. 1).

20 Acrp30 is a circulating protein that is produced by the adipose tissue (WO 96/39429 which is hereby incorporated herein in its entirety including any figures, tables, or drawings). Acrp30 mRNA levels are significantly reduced in adipose tissue from obese *ob/ob* mice. Acrp30 expression is greatly increased during adipocyte differentiation and appears to be regulated by insulin. In addition, Acrp30 has been shown to reduce postprandial blood lipid levels, stimulate weight loss, and reduce plasma triglycerides in normal mice, and to promote the reduction of food intake in genetically obese mice (PCT Application No. 99/07736 which is hereby incorporated herein in its entirety including any
25 figures, tables, or drawings).

30 Studies performed using just the globular region of Acrp30, termed globular Acrp30 (gAcrp30) showed that it effectively reduced body weight in mice on a high-fat cafeteria diet, as well as decreased fatty acid levels, decreased triglyceride levels, and improved glucose turnover in post-prandial tests in mice (See US Provisional Application No's 60/176,228 and 60/198,087, both of which are hereby incorporated herein by reference including any drawings, figures, or tables).

SUMMARY OF THE INVENTION

35 Globular OBG3 (gOBG3) has previously been linked with obesity, both in a murine model where treatment with gAcrp30 was shown to decrease body weight in mice fed a high fat diet, and in human subjects where an association between some Apm1 single nucleotide polymorphisms (SNPs)

and obesity was documented. The instant invention is drawn *inter alia* to the unexpected effects of gOBG3 on muscle cells, including increasing the oxidation of free fatty acid in muscle cells as well as accelerating muscle re-orientation/re-organization and differentiation.

5 In a first aspect, the invention features methods of accelerating muscle cell differentiation, comprising contacting muscle cells with gOBG3 thereby accelerating differentiation of the muscle cells. Preferably the contacting is performed under conditions such that gOBG3 binds to the muscle cells. Preferably the muscle cells are present in an individual. Preferably, gOBG3 is present in a pharmaceutical composition. The pharmaceutical composition preferably further comprises a pharmaceutically acceptable diluent. gOBG3 can be provided as a polypeptide or as a
10 polynucleotide encoding gOBG3. Preferably, gOBG3 is gApm1.

In a second aspect, the invention features methods of accelerating muscle cell reorganization, comprising contacting muscle cells with gOBG3 thereby accelerating reorganization of the muscle cells. Preferably the muscle cells are present in an individual. Preferably the contacting is under conditions such that gOBG3 binds to the muscle cells. Preferably, gOBG3 is
15 present in a pharmaceutical composition. The pharmaceutical composition preferably further comprises a pharmaceutically acceptable diluent. gOBG3 can be provided as a polypeptide or as a polynucleotide encoding gOBG3. Preferably, gOBG3 is gApm1.

In a third aspect, the invention features methods of accelerating muscle repair, comprising contacting muscle cells with gOBG3 thereby accelerating reorganization and differentiation of the
20 muscle cells. Preferably the contacting is under conditions such that gOBG3 binds to the muscle cells. Preferably the muscle cells are present in an individual. Preferably, gOBG3 is present in a pharmaceutical composition. The pharmaceutical composition preferably further comprises a pharmaceutically acceptable diluent. gOBG3 can be provided as a polypeptide or as a polynucleotide encoding gOBG3. Preferably, gOBG3 is gApm1.

25 In a fourth aspect, the invention features methods of increasing muscle mass in an individual, comprising contacting muscle cells in the individual with gOBG3 thereby accelerating the reorganization and differentiation of the muscle cells and increasing the muscle mass of the individual. Preferably said contacting is under conditions wherein gOBG3 binds to muscle cells. Preferably, gOBG3 is present in a pharmaceutical composition. The pharmaceutical composition
30 preferably further comprises a pharmaceutically acceptable diluent. gOBG3 can be provided as a polypeptide or as a polynucleotide encoding gOBG3. Preferably, gOBG3 is gApm1.

In a fifth aspect, the invention features methods of treating muscle cell disorders in an individual, comprising contacting muscle cells in the individual with gOBG3 thereby accelerating the reorganization and differentiation of the muscle cells, and thereby treating the muscle cell
35 disorders. Preferably the contacting is under conditions wherein gOBG3 binds to muscle cells. In preferred embodiments, the muscle cell disorders are selected from the group consisting of muscle-

related eye diseases and disorders, muscle-related recovery after injuries, muscle-related recovery after surgery, muscle-related disorders of aging, muscle atrophy and muscular dystrophy. Preferably, gOBG3 is present in a pharmaceutical composition. The pharmaceutical composition preferably further comprises a pharmaceutically acceptable diluent. gOBG3 can be provided as a polypeptide or as a polynucleotide encoding gOBG3. Preferably, gOBG3 is gApm1.

Definitions

Before describing the invention in greater detail, the following definitions are set forth to illustrate and define the meaning and scope of the terms used to describe the invention herein.

As used herein, the term "OBG3" refers to any member of the family of homologous proteins that includes Apm1, the human homologue, as well as Acrp30 or AdipoQ, the mouse homologue. These proteins and the polynucleotides encoding them are described in detail in WO 96/39429 and US Patent applications 60/176,228 and 60/198,087, all of which are hereby incorporated by reference herein in their entirety including any drawings, figures, or tables.

The term "polypeptide" refers to a polymer of amino acids without regard to the length of the polymer. Thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term also does not specify or exclude post-expression modifications of polypeptides. For example, polypeptides which include the covalent attachment of glycosyl groups, acetyl groups, phosphate groups, lipid groups and the like are expressly encompassed by the term polypeptide. Also included within the definition are polypeptides which contain one or more analogs of an amino acid (including, for example, non-naturally occurring amino acids, amino acids which only occur naturally in an unrelated biological system, modified amino acids from mammalian systems etc.), polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring.

As used herein, the term "non-human animal" refers to any non-human vertebrate, birds and more usually mammals, preferably primates, farm animals such as swine, goats, sheep, donkeys, and horses, rabbits or rodents, more preferably rats or mice. As used herein, the term "animal" is used to refer to any vertebrate, preferably a mammal. Both the terms "animal" and "mammal" expressly embrace human subjects unless preceded with the term "non-human".

The term "individual" as used herein refers to a mammal, including animals, preferably mice, rats, dogs, cattle, sheep, or primates, most preferably humans that perceives a need (or for whom a need is perceived) to accelerate muscle cell differentiation, to accelerate muscle cell reorganization, or to increase muscle mass. "Perceives a need" does not necessarily refer to a clinical need, but may simply be the result of an aesthetic desire to increase muscle mass, or to facilitate athletic training. An individual can also be a "patient".

The term "patient" as used herein refers to a mammal, including animals, preferably mice, rats, dogs, cattle, sheep, or primates, most preferably humans that are in need of treatment. The term "in need of treatment" as used herein refers to a judgment made by a medical care-provider such as a physician, nurse practitioner, nurse or the like, that a patient could benefit from or requires treatment. This judgement is made based on a variety of factors that are in the realm of the medical care-provider's expertise, but that include the knowledge that the patient is ill, or will be ill, as the result of a condition that is treatable by the compounds of the invention.

The term "accelerating muscle cell differentiation" as used herein refers to the ability of compounds of the invention to decrease the number of hours muscle cell differentiation requires in the absence of the compound (or alternatively stated, to increase the process of differentiation). This decrease in hours is at least 5 hours, preferably at least 10 hours, more preferably at least 20 hours. Alternatively, accelerating muscle cell differentiation can refer to the ability of compounds of the invention to increase the numbers of muscle cells undergoing differentiation at a given time, compared to the number in the absence of the compound, or to decrease the amount of apoptosis of muscle cells compared with the amount of apoptosis at a given time in the absence of the compound. Muscle cell differentiation can be determined by visual inspection (Example 1) or through markers of muscle cell differentiation known in the art, some of which are described in Shimokawa et al (1998) (Biochem Biophys Res Commun 246:287-292; hereby incorporated herein by reference in its entirety including drawings, figures, and tables; Example 6).

The term "accelerating muscle cell reorganization" as used herein refers to the ability of compounds of the invention to increase muscle cell reorganization in the presence of the compound as compared to in its absence. Alternatively, accelerating muscle cell reorganization can refer to the ability of compounds of the invention to increase the numbers of muscle cells undergoing reorganization at a given time, compared to the number in the absence of the compound. Muscle cell reorganization can be determined by visual inspection (Example 1).

The term "increasing muscle mass" as used herein refers to the ability of compounds of the invention to increase the number of differentiated muscle cells as compared to the numbers in the absence of the compound. Muscle cell differentiation can be determined by visual inspection (Example 1) or through markers of muscle cell differentiation known in the art, some of which are described in Shimokawa et al (1998). Increases in muscle mass can also be determined by measurements of the overall size of a muscle or the strength of a muscle using techniques well-known in the art.

The term "muscle cell disorders" as used herein refers to disorders where there is a loss of muscle mass, or muscle strength. These would include muscle atrophy or muscle wasting as the result of disease or trauma, or malnutrition, for example. In addition, disorders such as muscle-related eye diseases and disorders, muscle-related recovery after injuries, muscle-related recovery after

surgery, muscle-related disorders of aging, and muscular dystrophy are specifically envisioned. Whereas some of these disorders would require a medical care provider's diagnosis (muscular dystrophy for example), others would simply require an individual to detect the change and desire the treatment (some forms of muscle wasting or muscle atrophy).

5 The terms "percentage of sequence identity" and "percentage homology" are used interchangeably herein to refer to comparisons among polynucleotides and polypeptides, and are determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise
10 additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity. Homology is evaluated using any of the variety of
15 sequence comparison algorithms and programs known in the art. Such algorithms and programs include, but are by no means limited to, TBLASTN, BLASTP, FASTA, TFASTA, and CLUSTALW (Pearson and Lipman, 1988; Altschul et al., 1990; Thompson et al., 1994; Higgins et al., 1996; Altschul et al., 1990; Altschul et al., 1993). In a particularly preferred embodiment, protein and nucleic acid sequence homologies are evaluated using the Basic Local Alignment Search Tool
20 ("BLAST") which is well known in the art (see, e.g., Karlin and Altschul, 1990; Altschul et al., 1990; 1993, 1997). In particular, five specific BLAST programs are used to perform the following task:

- (1) BLASTP and BLAST3 compare an amino acid query sequence against a protein sequence database;
- 25 (2) BLASTN compares a nucleotide query sequence against a nucleotide sequence database;
- (3) BLASTX compares the six-frame conceptual translation products of a query nucleotide sequence (both strands) against a protein sequence database;
- (4) TBLASTN compares a query protein sequence against a nucleotide sequence database
30 translated in all six reading frames (both strands); and
- (5) TBLASTX compares the six-frame translations of a nucleotide query sequence against the six-frame translations of a nucleotide sequence database.

The BLAST programs identify homologous sequences by identifying similar segments, which are referred to herein as "high-scoring segment pairs," between a query amino or nucleic acid
35 sequence and a test sequence which is preferably obtained from a protein or nucleic acid sequence database. High-scoring segment pairs are preferably identified (i.e., aligned) by means of a scoring

matrix, many of which are known in the art. Preferably, the scoring matrix used is the BLOSUM62 matrix (Gonnet et al., 1992; Henikoff and Henikoff, 1993). Less preferably, the PAM or PAM250 matrices may also be used (see, e.g., Schwartz and Dayhoff, eds., 1978). The BLAST programs evaluate the statistical significance of all high-scoring segment pairs identified, and preferably
5 selects those segments which satisfy a user-specified threshold of significance, such as a user-specified percent homology. Preferably, the statistical significance of a high-scoring segment pair is evaluated using the statistical significance formula of Karlin (see, e.g., Karlin and Altschul, 1990).

By way of example and not limitation, procedures using conditions of high stringency are as follows: Prehybridization of filters containing DNA is carried out for 8 h to overnight at 65 °C in
10 buffer composed of 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 µg/mL denatured salmon sperm DNA. Filters are hybridized for 48 h at 65 °C, the preferred hybridization temperature, in prehybridization mixture containing 100 µg/mL denatured salmon sperm DNA and 5-20 X 10⁶ cpm of ³²P-labeled probe. Alternatively, the hybridization step can be performed at 65 °C in the presence of SSC buffer, 1 x SSC corresponding
15 to 0.15M NaCl and 0.05 M Na citrate. Subsequently, filter washes can be done at 37 °C for 1 h in a solution containing 2 x SSC, 0.01% PVP, 0.01% Ficoll, and 0.01% BSA, followed by a wash in 0.1 X SSC at 50 °C for 45 min. Alternatively, filter washes can be performed in a solution containing 2 x SSC and 0.1% SDS, or 0.5 x SSC and 0.1% SDS, or 0.1 x SSC and 0.1% SDS at 68 °C for 15 minute intervals. Following the wash steps, the hybridized probes are detectable by
20 autoradiography. Other conditions of high stringency may also be used and are well known in the art (Sambrook et al., 1989 and Ausubel et al., 1989, both of which are hereby incorporated herein in their entirety including any drawings, figures, or tables). These hybridization conditions are suitable for a nucleic acid molecule of about 20 nucleotides in length. The hybridization conditions described can be adapted according to the length of the desired nucleic acid, following techniques
25 well known to the one skilled in the art. The suitable hybridization conditions may, for example, be adapted according to Hames and Higgins (1985) or Sambrook et al. (1989).

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows an alignment of the sequences of the human (APM1), and mouse (adipoQ
30 and acrp30) OBG3 proteins.

Figure 2 shows a schematic drawing of the protein structure of APM1. The putative signal sequence at the N-terminus (AA 1-15), the unique region (AA 16-42), the collagen region (AA 43-107), and the globular region (AA 108-244) at the carboxy terminus are shown. Two protease cleavage sites at AA 100 and AA 131 are also shown.

35 Figure 3 shows a graphical representation of the effect of the addition of gACRP30 on the oleate oxidation of differentiated C2C12 cells.

Figure 4 shows a graphical representation of the effect of gAcrp30 treatment on fatty acid metabolism in muscle isolated from mice. EDL and Soleus muscles were isolated from both legs of each mouse (n=18). One muscle of each pair was incubated in medium with 2.5 µg/mL gAcrp30 (dark gray) and one in medium without gAcrp30 (control – light gray).

Figures 5A and 5B show graphical representations of the effect of gAcrp30 treatment on triglyceride content of muscle and liver isolated from mice. Mice that had received 25 µg of gAcrp30 twice daily showed significantly higher (p=0.002) muscle triglyceride content (Fig. 5A) than those receiving saline (control: light gray; gAcrp30: dark gray). This contrasted with a lack of increase in liver triglycerides (Fig. 5B).

Figures 6A and 6B show SDS-PAGE separations of the purification of Acrp30 and gAcrp30 (6A) and a cleavage product of apm1 (6B). Fig. 6A, Lane II shows the complete form of Acrp30 purified by FPLC. Lane I shows the proteolytic cleavage product gAcrp30. Fig. 6B shows a cleavage product of apm-1 after immunoprecipitation followed by Western blotting. The apparent molecular weight of this truncated form is 27kDa, corresponding to about 70% of the complete form of apm-1 (Lane IV). This truncated form was not detectable when a second anti-serum, specific for the human non-homologous region (HDQETTTQGPVLLPLPKGA) of the protein was used for immunoprecipitation (Lane V) and the same anti-globular head antiserum for detection. A preimmune serum of the same animal did not detect any protein; a dimer of apm-1 was seen with both specific antibodies (apparent MW 74kDa).

DETAILED DESCRIPTION OF THE INVENTION

Globular OBG3 has previously been linked with obesity, both in a murine model where treatment with globular Acrp30 (gAcrp30) was shown to decrease body weight in mice fed a high fat diet, and in human subjects where an association between some Apm1 single nucleotide polymorphisms (SNPs) and obesity was documented. In the instant application, the inventors have shown *inter alia* that at least some of the effects of gOBG3 are directed toward muscle cells. These effects include increasing the oxidation of free fatty acid in muscle cells as well as accelerating muscle re-organization and differentiation. Although the oxidation of free fatty acid in muscle cells likely is linked to the weight loss previously observed, it also seems to be linked to the acceleration of muscle cell reorganization and differentiation.

The effect of gAcrp30 on muscle cells was assessed using the murine, skeletal muscle cell line C2C12, and in *ex vivo* experiments on muscle cells excised from mice. The C2C12 cells were originally isolated from normal C3H mouse thigh muscle 72 hours after the muscle was crushed to increase the yield of mononucleated myogenic cells and designated C2 cells (Yaffe & Saxel (1977) Nature 270:725-727). The C2C12 cell line is a sub-clone of the C2 cell line selected for its ability to differentiate rapidly and to produce extensive contracting myotubes expressing characteristic muscle

proteins (Blau et al (1985) Science 230:758-766). Thus, it appears that the C2C12 line is probably a clonal derivative of the satellite muscle cells present in muscle tissue that can replicate and differentiate to form additional muscle fibers.

Differentiation of C2C12 myocytes is induced when cultures are shifted to medium containing low concentrations of mitogens (Wang & Walsh (1996) Science 273:359-361). During this process myoblasts withdraw permanently from the cell cycle, express muscle specific structural proteins, and fuse into multinucleated myotubes (Davis et al (1987). Extensive cell death is also observed in cultures of C2C12 cells exposed to differentiation medium containing 2% horse serum beginning at 24 hours and reaching a maximum of 20-30% of cells at 48 hours (Wang & Walsh (1996)). After 72 to 96 hours myotubes become abundant and cell death is diminished.

The process of differentiation of the C2C12 cell line is a good model for studies for treatments of damaged muscle tissue following muscle injuries associated with strains and sprains, and tears normally encountered in daily life, as well as during intense athletic training, or as the result of accidental injury, or surgery. In all these instances, the muscle tissue is damaged, needs to re-orient/reorganize, and to grow new muscle fibers. Thus, treatments that enhance the reorientation/reorganization of the cultured muscle cells and their differentiation into muscle fibers should also be useful for accelerating the healing of muscle tissue following injury, as well as for accelerating the augmentation or strengthening of muscle cells during physical therapy, or athletic training.

Globular Acp30 was found to induce pronounced re-orientation/re-organization of undifferentiated C2C12 muscle cells, as well as to accelerate the process of differentiation into muscle fibers. Further, there were indications of a decrease in apoptosis of muscle cells during the differentiation process, since the numbers of cells compared to the cells not treated with gOBG3 increased. Treatment of differentiated C2C12 cells with gAcp30 also caused an increase in fatty acid metabolism, since oleate oxidation was increased approximately 40% (Fig. 3). A significant increase in oleate oxidation was also seen in *ex vivo* experiments with isolated mouse EDL and soleus muscles (Fig. 4), indicating the strength of the C2C12 cell line as a model system. A concurrent significant increase in triglyceride concentration was also observed in the *ex vivo* muscle.

Although not wishing to be limited to one hypothesis, the inventors believe that the increase in metabolism of free fatty acids that results from the addition of gAcp30 may be involved in the acceleration of muscle cell reorientation/reorganization and differentiation. It is possible that the increase in fatty acid oxidation provides nutrients or energy that are involved in the process, or simply sends a signal. Whatever the exact mechanism involved, it is clear that treatment with gOBG3 has dramatic effects on muscle cells both *in vitro* and *in vivo*, for re-organization, differentiation and preventing apoptosis. Thus, it should be useful for treatment of muscle disorders where additional muscle tissue is desired and potentially where prevention of muscle cell death is

needed. Examples of such disorders include, but are not limited to, muscle wasting, muscle atrophy, or muscular dystrophy. Augmenting muscle differentiation and growth should be ameliorative to the symptoms, if not curative of the disease. Recruitment of more muscle cells, alignment, and differentiation of more muscle fibers should have a positive effect, and might at least prolong the useful life of the muscles of patients afflicted with muscular dystrophy.

PREFERRED EMBODIMENTS OF THE INVENTION

I. Muscle-related Uses of gOBG3

Methods of Accelerating Muscle Repair:

The inventors have shown that treatment of muscle cells with gOBG3 leads to their re-organization and differentiation, as well as increased free fatty acid oxidation, processes believed to be important in muscle repair. The muscle cell line used is one created as the result of trauma to skeletal muscle cells and is therefore thought to be a good model for studying muscle repair.

Treatments that accelerate muscle cell re-organization, muscle cell differentiation, and/or muscle cell repair would be useful following any kind of muscle injury, including, but not limited to trauma, either accidental, or the result of surgery, or over-exercising. Trauma to muscles can result from blows, tears, cuts, strains, etc. gOBG3 variants and fragments, as well as agonists and antagonists of gOBG3, can be tested for their activity and thus their potential for use as treatments for the acceleration of muscle repair using the assays described in the Examples (particularly Examples 1, 2, 4, 6, 7) or in other assays known to those in the art.

Methods of Increasing Muscle Mass:

In addition to muscle repair, the results of treatment of muscle cells with gOBG3 suggest that gOBG3 treatment could also be useful for increasing muscle mass and/or increasing muscle strength and/or muscle endurance. Increasing body mass (for aesthetic or sports-related reasons, for example) also involves the recruitment and development of new muscle cells, which gAcrp30 has been shown to promote in the experiments with C2C12 cells. Further, increased free fatty acid oxidation should also be useful in any kind of endurance or other activities leading to muscle fatigue, since free fatty acids are a better source of energy and generally less easily utilized than glucose stores. To some extent, gOBG3 would be expected to function similarly to the anabolic steroid-type drugs currently used by athletes. gOBG3 variants and fragments, as well as agonists and antagonists of gOBG3, can be tested for their activity and thus their potential for use for increasing muscle mass, strength and/or endurance using the assays described in the Examples (particularly Examples 1, 2, 4, 6, 7) or assays known to those in the art.

Methods of Treatment of Muscle Disorders:

For similar reasons as those that suggest gOBG3 would be useful for accelerating muscle repair and increasing muscle mass, and additionally because it appears that gOBG3 may be useful in preventing apoptosis, gOBG3 should also be useful for treating muscle cell disorders. The muscle cell disorders contemplated are those that would improve, or whose symptoms would be ameliorated by treatment with gOBG3. These would include disorders in which the cells need to be strengthened (improve utilisation of free fatty acid), or the amount of muscle fibers increased (increased differentiation of muscle cells). For example, gOBG3 would be expected to be useful for treating the muscle cell disorders muscle atrophy, muscle wasting, and muscular dystrophy. Treatment with gOBG3 is expected to ameliorate some symptoms of these diseases by increasing the strength of the existing muscle cells by increasing their use of free fatty acids, and by increasing the differentiation of additional muscle cells, as well as by preventing the apoptosis of existing muscle cells. For instance, even though in muscular dystrophy the existing muscles are abnormal and their use is gradually lost, it is thought that gOBG3 should be able to increase the useful life of the muscles. gOBG3 variants and fragments, as well as agonists and antagonists of gOBG3, can be tested for their activity and thus their potential for use as treatments for muscle disorders using the assays described in the Examples (particularly Examples 1, 2, 4-8) or in other assays known to those in the art.

II. Globular OBG3 Polypeptides

Globular OBG3 polypeptides are used in the methods of treating muscle cells of the instant invention. As used herein, the term "gOBG3" refers to the globular portion of any member of the family of homologous proteins that includes Apm1, the human homologue, as well as Acrp30 or AdipoQ, the mouse homologue. Globular OBG3 polypeptides have previously been described in detail in US Provisional patent application Nos 60/176,228 and 60/198,087, hereby incorporated by reference herein in their entirety including figures, drawings, or tables. As used herein, unless specifically limited, the term is meant to include modified gOBG3 polypeptide sequences, including variants, fragments, analogs and derivatives of the gOBG3 polypeptides as described previously.

For the purposes of this invention, however, useful gOBG3 polypeptides are those that retain any one or more of the desired activities described herein, including but not limited to the effects on muscle cells that are the subject of the instant invention. These include accelerating re-orientation and differentiation of muscle cells as well as increasing free fatty acid oxidation, and preventing apoptosis. Variants, fragments, analogs and derivatives of these polypeptide sequences can be assayed for their retention of the desired activities using any of the methods/tests described in Examples 1,2, and 4-8 or any comparable assays.

Globular OBG3 is the portion of intact OBG3 that does not include the collagen-like tail, or that contains few enough of the collagen residues such that the peptides do not assemble, or if they assemble this does not inhibit their activity. Preferably, this is fewer than 6 collagen residues, fewer

than 4, fewer than 2, or more preferably no collagen residues. By "collagen residues" as used herein is meant the amino acids glycine, X, Y, where X and Y can be any amino acid. The collagen-like region of OBG3 is shown in Fig. 2 for APM1.

The term "activity" as used herein refers to a measurable result of the interaction of molecules. For example, a preferred gOBG3 activity is to accelerate re-orientation of muscle cells, accelerate differentiation of muscle cells, and/or increase free fatty acid oxidation of muscle cells. Representative assays to test for these functions are provided in Examples 1,2, and 4-9. However, these examples are provided for explanation, not limitation. Those with skill in the art would be able to design other experiments to test for the same retained activity.

The term the "same retained activity" as used herein refers to the ability of a variant, fragment, analog or derivative of gOBG3 to have the same activity as is demonstrated in Examples 1,2, and 4-8 and claimed in the instant invention. The variant, fragment, analog or derivative of gOBG3 does not necessarily have to retain all of the activities described herein for gOBG3's action on muscle cells, unless specified, but preferably retains at least one of the activities.

The "same activity" also relates to the amount of a given activity observed. In the instant application this refers to the amount of oleate oxidation, or amount of acceleration of differentiation, or amount of prevention of apoptosis, for example. Preferably, the "same" activity as it relates to amount, means within 10% of the previously observed amount, but it can include a difference of 20% or 30% or even 50%. However, this is not meant to limit the use of more effective gOBG3 polypeptides.

"More effective" gOBG3 polypeptides include those with an increased activity compared with the gAcrp30 polypeptide used in experiments described herein. The term "increased" as used herein refers to the ability of gOBG3 polypeptides to increase an activity in some measurable way as compared to an appropriate control. As a result of the presence of a gOBG3 variant, the levels of fatty acid oxidation, or muscle cell differentiation might increase, or the amount of apoptosis might decrease, for example, as compared to appropriate controls, typically the presence of gAcrp30 used in the experiments described herein. Preferably, an increase in activity is at least 25%, more preferably at least 50%, most preferably at least 100%.

"More effective" gOBG3 polypeptides may also include those that lack or have a decreased amount of one activity compared with the gAcrp30 polypeptide used in experiments described herein, and an increased amount of another activity compared with the gAcrp30 polypeptide used in experiments described herein. The term "decreased" as used herein refers to the ability of gOBG3 polypeptides to decrease an activity in some measurable way as compared to an appropriate control, such as the gAcrp30 polypeptide used in experiments described herein. As a result of the presence of a gOBG3 variant, fatty acid oxidation might decrease, for example, as compared to controls in the presence of the gAcrp30 polypeptide used in experiments described herein. Preferably, a decrease in

activity is at least 25%, more preferably at least 50%, most preferably at least 100%. The term "lack" as used herein refers to an inability to detect an activity using the methods described herein, or similar methods. A gOBG3 variant could be thought to "lack" activity even though an increase of 5 or 10 or 15% of an effect is observed compared with an assay performed in its absence.

5 Finally, "more effective" gOBG3 polypeptides may also include those that lack or have a decreased amount of one activity or all activities compared with the gAcrp30 polypeptide used in experiments described herein, but an increased amount of these activities *in vivo* as compared with the gAcrp30 polypeptide used in experiments described herein.

Preferred embodiments of the invention feature gOBG3 polypeptide that consists of the
10 sequence of the globular region shown in Figure 1, or variants, fragments, analogs, or derivatives thereof. Preferable embodiments include amino acids 108-244 of SEQ ID NO:6 or 111-247 of SEQ ID Nos 2 and 4. Alternative preferable embodiments include amino acids 104 to 247 of the OBG3 proteins described in Figure 1.

In other preferred embodiments, the invention features a gOBG3 polypeptide comprising at
15 least 115, but not more than 175 contiguous amino acids of any one of the gOBG3 polypeptide sequences set forth in Figure 1, wherein no more than 12 of said at least 115 and no more than 175 contiguous amino acids are present in the collagen-like region of OBG3. Preferably, the gOBG3 polypeptide comprises at least 125, but not more than 165, or at least 135, but not more than 155, and no more than 9 are in the collagen-like region; more preferably at least 125 but not more than
20 165, or 135 but not more than 155, and no more than 6 are in the collagen-like region; or at least 140 and not more than 150, and no more than 3 are present in the collagen-like region. Preferably the gOBG3 polypeptide is human or mouse, but most preferably human.

Variant gOBG3 polypeptides of the invention may be 1) ones in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue and such
25 substituted amino acid residue may or may not be one encoded by the genetic code, or 2) ones in which one or more of the amino acid residues includes a substituent group, or 3) ones in which a modified gOBG3 polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or 4) ones in which the additional amino acids are fused to modify a gOBG3 polypeptide, such as a leader or secretory sequence or a
30 sequence which is employed for purification of the modified gOBG3 polypeptide or a pre-protein sequence. Such variants are deemed to be within the scope of those skilled in the art. The retention of the desired activity (and thus desired gOBG3 polypeptides) can be determined using the assays described in Examples 1, 2, 4-8 or other assays that achieve the same result.

Amino acid changes present in a variant polypeptide may be non-conservative amino acid
35 changes but more preferably are conservative amino acid changes. In cases where there are one or more amino acid changes, preferred gOBG3 polypeptides include those that retain the same

activities and activity levels as the reference gOBG3 polypeptide sequence, as well as those where the level of one or more activities is increased. Assays for determining gOBG3 polypeptide activities of the invention are described herein in the Examples (1, 2, 4-8) in more detail, but include accelerating muscle differentiation, muscle oleate oxidation, and decreasing muscle apoptosis, both
5 *in vitro* and *in vivo*.

In preferred embodiments, the invention features a variant of a gOBG3 polypeptide that is at least 75% identical to gOBG3 polypeptide sequences selected from the group consisting of 101-244, 108-244, and 132-244 of SEQ ID NO:6, or 104-247, 111-247, and 135-247 of SEQ ID Nos 2 or 4. Preferably, the amino acid sequence is at least 85% identical, more preferably 90% identical, most
10 preferably 95% identical and optionally 100% identical. Preferably the sequence is human or mouse, and most preferably human.

In yet other preferred embodiments, the invention features a variant of a gOBG3 polypeptide that comprises (or consists of) a 143 contiguous amino acid sequence, wherein at least 100 of the 143 amino acids are identical to amino acids 101-244 of SEQ ID NO:6 or 104-247 of SEQ ID Nos 2
15 or 4. Preferably, at least 113 of the 143 amino acids are identical, more preferably 127 of the 143 are identical, even more preferably 134 of the 143 are identical, and most preferably all of the amino acids are identical. Preferably the sequence is human or mouse, and most preferably human.

In yet other preferred embodiments, the invention features a variant of a gOBG3 polypeptide that comprises (or consists of) a 137 contiguous amino acid sequence, wherein at least 100 of the
20 137 amino acids are identical to amino acids 108-244 of SEQ ID NO:6 or 111-247 of SEQ ID Nos 2 or 4. Preferably, at least 113 of the 137 amino acids are identical, more preferably 127 of the 137 are identical, even more preferably 134 of the 137 are identical, and most preferably all of the amino acids are identical. Preferably the sequence is human or mouse, and most preferably human.

In yet other preferred embodiments, the invention features a variant of a gOBG3 polypeptide
25 that comprises (or consists of) a 113 contiguous amino acid sequence, wherein at least 80 of the 113 amino acids are identical to amino acids 132-244 of SEQ ID NO:6 or 135-247 of SEQ ID Nos 2 or 4. Preferably, at least 90 of the 113 amino acids are identical, more preferably 100 of the 113 are identical, even more preferably 110 of the 113 are identical, and most preferably all of the amino acids are identical. Preferably the sequence is human or mouse, and most preferably human.

30 A polypeptide fragment is a polypeptide having a sequence that is entirely the same as part, but not all, of a given polypeptide sequence, preferably a gOBG3 polypeptide and variants thereof. Such fragments may be "free-standing", *i.e.* not part of or fused to other polypeptides, or they may be comprised within a single larger non-OBG3 polypeptide of which they form a part or region. However, several fragments may be comprised within a single larger polypeptide. As representative
35 examples of gOBG3 polypeptide fragments of the invention, there may be mentioned those which have from about 5, 6, 7, 8, 9 or 10 to 15, 10 to 20, 15 to 40, 30 to 55, 40 to 70, 60 to 95, 80 to 130, or

90 to 144 amino acids long. Preferred are those fragments containing at least one amino acid substitution or deletion compared to a gOBG3 polypeptide.

III. Pharmaceutical Compositions

5 The gOBG3 polypeptides of the invention can be administered to a mammal, including a human patient, alone or in pharmaceutical compositions where they are mixed with suitable carriers or excipient(s). The pharmaceutical composition is then provided at a therapeutically or aesthetically effective dose. A therapeutically or aesthetically effective dose refers to that amount of gOBG3 sufficient to result in amelioration of symptoms of muscle-related disorders as determined by the methods described herein. A therapeutically or aesthetically effective dose can also refer to the amount of gOBG3 necessary for an increase in muscle mass or an increase in muscle strength or endurance in persons desiring this affect for aesthetic or athletic reasons alone. A therapeutically effective dosage of a gOBG3 polypeptide of the invention is that dosage that is adequate to promote muscle differentiation and/or increased free fatty acid oxidation, or decreased muscle cell apoptosis with continued or periodic use or administration. Techniques for formulation and administration of gOBG3 may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition.

Other diseases or disorders that gOBG3 could be used to treat or prevent include, but are not limited to, muscle atrophy, muscle wasting and muscular dystrophy.

20

Routes of Administration.

Suitable routes of administration include oral, rectal, transmucosal, or intestinal administration, parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal or intraocular injections. A particularly useful method of administering compounds for promoting weight loss involves surgical implantation, for example into the abdominal cavity of the recipient, of a device for delivering gOBG3 over an extended period of time. Sustained release formulations of the invented medicaments particularly are contemplated.

30 Composition/Formulation

Pharmaceutical compositions and medicaments for use in accordance with the present invention may be formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries. Proper formulation is dependent upon the route of administration chosen.

35 Certain of the medicaments described herein will include a pharmaceutically acceptable carrier and at least one polypeptide that is a gOBG3 polypeptide of the invention. For injection, the agents of

the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer such as a phosphate or bicarbonate buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

5 Pharmaceutical preparations that can be taken orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with fillers such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty
10 oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

For administration by inhalation, the compounds for use according to the present invention are
15 conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable gaseous propellant, *e.g.*, carbon dioxide. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, *e.g.*, gelatin, for use in an inhaler or insufflator, may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or
20 starch.

The compounds may be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, *e.g.*, in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in aqueous vehicles, and may contain formulatory
25 agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Aqueous suspensions may contain substances that increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents that increase the solubility of
30 the compounds to allow for the preparation of highly concentrated solutions.

Alternatively, the active ingredient may be in powder or lyophilized form for constitution with a suitable vehicle, such as sterile pyrogen-free water, before use.

In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for
35 example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an

emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various sustained release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days.

Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

The pharmaceutical compositions also may comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

Effective Dosage.

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve their intended purpose. More specifically, a therapeutically effective amount means an amount effective to prevent development of or to alleviate the existing symptoms of the subject being treated. Determination of the effective amounts is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. For example, a dose can be formulated in animal models to achieve a circulating concentration range that includes or encompasses a concentration point or range shown to increase leptin or lipoprotein uptake or binding in an *in vitro* system. Such information can be used to more accurately determine useful doses in humans.

A therapeutically effective dose refers to that amount of the compound that results in amelioration of symptoms in a patient. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50, (the dose lethal to 50% of the test population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD50 and ED50. Compounds that exhibit high therapeutic indices are preferred.

The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50, with little or no toxicity. The dosage may vary

within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See, e.g., Fingl et al., 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1).

5 Dosage amount and interval may be adjusted individually to provide plasma levels of the active compound which are sufficient to maintain the weight loss or prevention of weight gain effects. Dosages necessary to achieve these effects will depend on individual characteristics and route of administration.

10 Dosage intervals can also be determined using the value for the minimum effective concentration. Compounds should be administered using a regimen that maintains plasma levels above the minimum effective concentration for 10-90% of the time, preferably between 30-90%; and most preferably between 50-90%. In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

15 The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

A preferred dosage range for the amount of a gOBG3 polypeptide of the invention, that can be administered on a daily or regular basis to achieve desired results, including a reduction in levels of circulating plasma triglyceride-rich lipoproteins, range from 0.01 - 50 mg/kg body mass. A more preferred dosage range is from 0.02 - 25 mg/kg. A still more preferred dosage range is from 0.1 - 20 mg/kg, while the most preferred range is from 0.2 - 10 mg/kg. Of course, these daily dosages can be delivered or administered in small amounts periodically during the course of a day.

EXAMPLES

25 Other characteristics and advantages of the invention are described in the Examples. The following Examples are provided for illustrative purposes and not as a means of limitation. One of ordinary skill in the art would be able to design equivalent assays and methods based on the disclosure herein, all of which form part of the instant invention.

30 Throughout this application, various publications, patents and published patent applications are cited. The disclosures of these publications, patents and published patent specifications referenced in this application are hereby incorporated by reference into the present disclosure.

EXAMPLE 1: Effect of gACRP30 Treatment of Muscle Cells on Muscle Differentiation

35 C2C12 cells (murine skeletal muscle cell line; ATCC CRL 1772, Rockville, MD) are seeded sparsely (about 15-20%) in complete DMEM (w/glutamine, pen/strep, etc) + 10% FCS. Two days later they become 80-90% confluent. At this time, the media is changed to DMEM+2% horse serum

to allow differentiation. The media is changed daily. Abundant myotube formation occurs after 3-4 days of being in 2% horse serum.

To test the effect of the presence of gACRP30 on muscle differentiation, gACRP30 (1 to 2 $\mu\text{g/mL}$) was added the day after seeding when the cells are still in DMEM w/ 10% FCS. Two days after plating the cells (one day after gACRP30 was first added), at about 80-90% confluency, the media was changed to DMEM+2% horse serum plus gACRP30.

The results show that the addition of gACRP30 causes the cells to begin organizing within one day after its addition. In contrast to the random orientation of the cells not treated with gACRP30, those treated with gACRP30 aligned themselves in relation to each other. In addition, differentiation progressed more rapidly in the presence of gACRP30; after only 2 days of gACRP30 treatment maximal or near maximal differentiation was observed in contrast to the 3 to 4 days needed in its absence.

EXAMPLE 2: Effect of gACRP30 on Muscle Cell Fatty Acid Oxidation

C2C12 cells were differentiated in the presence or absence of 2 $\mu\text{g/mL}$ gACRP30 for 4 days. On day 4, oleate oxidation rates were determined by measuring conversion of 1- ^{14}C -oleate (0.2 mM) to $^{14}\text{CO}_2$ for 90 min. C2C12 cells differentiated in the presence of gACRP30 undergo 40% more oleate oxidation than controls differentiated in the absence of oleate (Fig. 3).

The effect of gACRP30 on the rate of oleate oxidation was compared in differentiated C2C12 cells (murine skeletal muscle cells; ATCC, Manassas, VA CRL-1772) and in a hepatocyte cell line (Hepa1-6; ATCC, Manassas, VA CRL-1830). Cultured cells were maintained according to manufacturer's instructions.

The oleate oxidation assay was performed as previously described (Muoio et al (1999) Biochem J 338:783-791). Briefly, nearly confluent myocytes were kept in low serum differentiation media (DMEM, 2.5% Horse serum) for 4 days, at which time formation of myotubes became maximal. Hepatocytes were kept in the same DMEM medium supplemented with 10% FCS for 2 days. One hour prior to the experiment the media was removed and 1 mL of preincubation media (MEM, 2.5% Horse serum, 3 mM glucose, 4 mM Glutamine, 25 mM Hepes, 1% FFA free BSA, 0.25 mM Oleate, 5 $\mu\text{g/mL}$ gentamycin) was added. At the start of the oxidation experiment, ^{14}C -Oleic acid (1 $\mu\text{Ci/mL}$, American Radiolabeled Chemical Inc., St. Louis, MO) was added and cells were incubated for 90 min at 37 $^{\circ}\text{C}$ in the absence/presence of 2.5 $\mu\text{g/mL}$ gACRP30. After the incubation period, 0.75 mL of the media was removed and was assayed for ^{14}C -oxidation products as described for the muscle FFA oxidation experiment.

Oleate oxidation in C2C12 cells determined over 90 min increased significantly (39%; $p = 0.036$, two-tailed t-Test) in cells treated with gACRP30. In contrast, no detectable increase in the rate of FFA oxidation was seen in hepatocytes incubated with gACRP30.

Triglyceride and Protein Analysis following Oleate Oxidation in cultured cells

Following the transfer of media for the oleate oxidation assay, cells were placed on ice. To determine triglyceride and protein content, cells were washed with 1 mL of 1x PBS to remove residual media. To each well, 300 μ L of cell dissociation solution (Sigma) was added. Cells were
 5 incubated at 37 °C for 10 min. Plates were tapped to loosen cells, and 0.5 mL of 1x PBS was added. The cell suspension was transferred to an eppendorf tube, each well was rinsed with an additional 0.5 mL of 1x PBS, and the rinse was transferred to appropriate eppendorf tube. Samples were centrifuged at 1000 rpm for 10 minutes at room temperature. The supernatant was discarded, and 750 μ L of 1x PBS/2% chaps was added to each cell pellet. The cell suspension was vortexed and
 10 was placed on ice for 1 hour. Samples were then centrifuged at 13000 rpm for 20 min at 4 °C. Supernatants were transferred to new tubes and frozen at -20 °C until analyzed.

The amount of triglycerides in each sample was determined using Sigma Diagnostics GPO-TRINDER enzymatic kit. The procedure outlined in the manual was adhered to with the following exceptions: the assay was performed in a 48 well plate, 350 μ L of sample volume was assayed, the
 15 control blank consisted of 350 μ L PBS/2% chaps, and the standard contained 10 μ L of the standard provided in the kit plus 690 μ L PBS/2% chaps. Analysis of the samples was performed on a Packard Spectra Count at a wavelength of 550 nm.

Protein analysis was carried out on 25 μ L of each supernatant sample using the BCA protein assay (Pierce) following manufacturer's instructions. Analysis of the samples was performed on a
 20 Packard Spectra Count at a wavelength of 550 nm.

Triglyceride production in both C2C12 and Hepa 1-6 cells did not change significantly in the absence/presence of Acrp30 and gAcrp30. The protein content of all cells analyzed was equivalent in the absence/presence of Acrp30 and gAcrp30.

25 EXAMPLE 3: Production of Recombinant Acrp30

An illustrative method for producing recombinant OBG3 is given below. Although the method describes the production of the mouse analog, Acrp30, a person of ordinary skill in the art would be able to use the guidance provided to produce other OBG3 analogs, including but not limited to, the human analog, Apm1.

30 Recombinant Acrp30 is cloned in pTRC His B (Invitrogen) between BamH1 and Xho1 and maintained in *E. coli* DH5- α . The sequence of the Acrp30 insert corresponds to Acrp 30 genbank U37222 bases 88 to 791, except in position 382 where in #3 G replaces A found in ACRP 30 (V instead of M). The corresponding nucleotide in AdipoQ U49915 is G as in clone #3. The amino acid V is also conserved in the human sequence Apm-1 D45371.

35

Culture:

Plate out bacteria in LB agar media containing 100 µg/mL ampicillin. Inoculate 1 colony into 5 mL media (no agar) at 37 °C overnight. Add 2 mL of this initial culture into 500 mL Erlenmeyer flasks containing 200 mL LB media and 100 µg/mL ampicillin. Incubate at 37 °C in an orbital shaker until the OD₆₀₀ = 0.2. Add IPTG to a final concentration of 1 mM (stock solution = 1 M). Incubate at 37 °C overnight.

Lysis:

Pellet the bacteria by centrifugation (Sorvall, 3500 rpm, 15 min, 4 °C) in a pre-weighed tube.

At 4 °C resuspend the pellet in 3 mL/g of lysis buffer

Add 40 µL/g PMSF 10 mM

Add 80 µL/g of lysozyme 10 mg/mL

Incubate 20 min on ice, shaking intermittently

Add 30 µL/g 10 % sodium deoxycholate

Incubate at 37 °C until the lysate is viscous

Freeze in liquid Nitrogen and thaw at 37 °C three times

Sonicate 2X, 30 sec, 25 % cycle, 2.5 power level

Centrifuge 30 min, 15000 rpm, 4 °C

Recover the supernatant

Note: The lysate can be stored frozen before or after the sonication step.

Batch Purification:

1. Pack 1 mL of Probond resin (Invitrogen; 1 mL = 2 mL suspended gel) into a 5 mL column. Wash with 5 mL PBS.

2. Apply 5 mL bacterial supernatant to the 1 mL of gel. (If volume is very high, use several small columns.)

3. Wash with 24 mL phosphate buffer, pH 7.8, followed by a wash with 24 mL phosphate buffer, pH 6.

4. Elute with imidazole buffer and collect fractions of 1 mL.

5. Analyze fractions by OD at 280 nm or by SDS-PAGE (12.5 %; dilution ½ in 2X sample buffer) under reducing conditions (100 °C, 5 min)

6. Pool the fractions containing protein (usually fraction numbers 2-4 for concentrations of 0.8 – 1 mg/mL and fractions 1, 5 and 6 for concentrations of 0.2 – 0.4 mg/mL).

7. Dialyze thoroughly against 1 X PBS, 24 mM ammonium bicarbonate or 50 mM Tris, pH 7.4 containing 250 mM NaCl. Concentrate by Speed-Vac if needed.

8. Analyze protein by the Lowry method.

9. Aliquot and store at -20°C .

Purification On Liquid Chromatography System

1. Pack 5 mL of Probond resin into a 5 mL column.
- 5 2. Wash with 4 bed volumes of phosphate buffer pH 7.8, 1 mL/min.
3. Inject 25 mL lysate (filtered on $0.45\ \mu$ or centrifuged at 3000 rpm, 30 min, 4°C , Beckman Allegra 6R) at 0.5 mL/min.
4. Wash with 4 bed volumes of phosphate buffer, pH 7.8 at 1 mL/min.
5. Wash with 12 bed volumes of phosphate buffer pH 5.5 at 1 mL/min.
- 10 6. Elute bound fraction with phosphate buffer, pH 5.5, containing 1 M imidazole at 1 mL/min.
7. Collect fractions, dialyze and analyze protein as described for batch purification, steps 7-9.

15 EXAMPLE 4: Generation of Globular Acrp30 by Enzymatic Cleavage

Incubate purified Acrp30 (obtained as described above or through equivalent method) with acetylated Trypsin-Type V-S from Bovine Pancreas (Sigma E.C. = 3.4.21.4) at 400 u/mg protein at 25°C for 10 min.

- 20 Stop reaction by running the sample over a Poly-Prep Column (Biorad 731-1550) at $+4^{\circ}\text{C}$ containing immobilized Trypsin inhibitor.

Collect 1.0 mL fractions. Determine protein concentration.

Pool the protein containing fractions and dialyze extensively against PBS using dialysis tubing with M.W. cutoff = 10,000 da.

- 25 Concentrate on Amicon YM-10 Centricon Filter (Millipore, M.W. cutoff = 10,000 da).
- Sterile filter.

Determine final protein concentration using Markwell's modified Lowry procedure (1981) or BCA protein assay (Pierce Chemical Co, Rockford, IL) and BSA as standard.

- 30 Check purity and efficiency of cleavage by SDS – PAGE analysis using a 4-20% gradient gel. The intact Acrp30 migrates as a single band at approx. 37 kda due to co-transcribed vector sequences attached to the histidine tag at the N-terminus of Acrp30. The cleaved Acrp30 forms a band at approx. 18 kda (gAcrp30). Additional degradation products, all smaller than 10 kda are also generated from the N-terminal region. These are separated from the desired 18 kda band by dialysis with semipermeable membranes with a MW cutoff of 10,000. The actual cleavage site using this method has been identified as the one after amino acid 103.

35 Example 4: Effect of ACRP30 on Oleate Oxidation in Isolated Muscle

Intact muscles were isolated from C57BL6/J mice and the oleate oxidation of the isolated muscle was measured (Clee et al (2000) J Lipid Res 41:521-531; Muoio et al (1999) Am J Physiol 276:E913-921). Oleate oxidation in isolated muscle was measured as previously described (Cuendet et al (1976) J Clin Invest 58:1078-1088; Le Marchand-Brustel (1978) Am J Physiol 234:E348-E358, hereby incorporated by reference herein in its entirety including any figures, drawings, or tables).

Two groups of C57BL6/J mice, age 5 and 33 weeks, were used. Mice were kept on a regular diet with free access to food and water. The day/night light cycle was kept at 12hr ON/12hr OFF. The older animals were injected twice daily with either saline or 25 µg of gACRP30 for 3-4 days. The younger group of animals was given a high fat meal (6g butter, 6g sunflower oil, 10 g nonfat dry milk, 10 g sucrose, 12 mL distilled water prepared fresh) by gavage (vol.=1% of body weight) at time 0 and was injected with saline or gACRP30 only at time 0 and again at 45 minutes. They were sacrificed at 180 minutes. The gACRP30 used was prepared as described in Example 3.

After cervical dislocation, soleus and EDL muscles were rapidly isolated from the hind limbs of the mice. The distal tendon of each muscle was tied to a piece of suture to facilitate transfer among different media. All incubations were carried out at 30 °C in 1.5 mL of Krebs-Ringer bicarbonate buffer (118.6 mM NaCl, 4.76 mM KCl, 1.19 mM KH₂PO₄, 1.19 mM MgSO₄, 2.54 mM CaCl₂, 10 mM Hepes, pH 7.4) supplemented with 4% bovine serum albumin, FFA free (fraction V, RIA grade, Sigma, Inc., St. Louis, MO) and 5 mM glucose (Sigma, Inc., St. Louis, MO). The concentration of oleate (Sigma, Inc., St. Louis, MO) throughout the experiment was 0.25 mM. All media were oxygenated (95% O₂; 5% CO₂) prior to incubation. The gas mixture was hydrated throughout the experiment by bubbling through a gas washer (Kontes Inc., Vineland, NJ).

Muscles were rinsed for 30 min in incubation media with oxygenation. The muscles were then transferred to fresh media (1.5 mL) and incubated at 30 °C in the presence of 1 µCi/mL [1-¹⁴C] oleic acid (ARC) and 1 µCi/mL [5-³H]-glucose (Amersham). The incubation vials containing this media were sealed with a rubber septum from which a center well carrying a piece of Whatman paper (1.5 cm x 11.5 cm) was suspended.

After an initial incubation period of 10 min with constant oxygenation, gas circulation was removed to close the system to the outside environment. The muscles were incubated for an additional 90 min at 30 °C. At the end of this period, 0.45 mL of Solvable (Packard Instruments, Meriden, CT) was injected onto the Whatman paper in the center well. The oxidation of oleate by the muscle was stopped at this time point by transferring the vial onto ice.

After 5 min, the muscle was removed from the medium, was cleaned of connective tissue, was weighed and was frozen for further analysis. An aliquot of 0.5 mL medium was removed, the vials were closed again and 1 mL of 35% perchloric acid was injected with a syringe into the media by piercing through the rubber septum with a needle. The CO₂ released from the acidified media was collected by the Solvable in the center well.

After a 90 min collection period at 30 °C, the Whatman paper was removed from the center well and placed in scintillation vials containing 15 mL of scintillation fluid (HionicFlour, Packard Instruments, Meriden, CT). The amount of ^{14}C radioactivity was quantitated by liquid scintillation counting. The rate of oleate oxidation was expressed as nmol oleate produced in 90 min/g muscle.

5 To test the effect of gACRP30 or ACRP30 on oleate oxidation, these proteins were added to the media at a final concentration of 2.5 $\mu\text{g/mL}$ and maintained in the media throughout the procedure. Two muscles of different oxidative capacity (soleus and extensor digitorum longus (EDL)) were tested (Fig. 4). EDL and Soleus muscles were isolated from both legs of normal C57BL/6J mice ($n=18$). One muscle of each pair was incubated in medium with 2.5 $\mu\text{g/mL}$ gAcrp30 (dark gray) and one in medium without gAcrp30 (control – light gray). This experimental
10 design allowed us to compare oleate oxidation in pairs of muscles isolated from the same animal. ^{14}C -Oleate oxidation was determined over 90 minutes. Incubation of EDL and soleus muscles for 90 minutes in medium containing 2.5 $\mu\text{g/mL}$ gAcrp30 leads to a statistically significant increase in oleate oxidation ($p<0.05$, paired, one-tailed, t-Test) or ($p=0.0041$, Repeated Measures Analysis of
15 Variance, Univariate Tests of Hypotheses for Within Subject Effects) in both muscle types.

Both muscle types showed a significant response to gAcrp30. The relative increase in FFA oxidation was 17% ($p=0.03$) and 10% ($p=0.04$) for EDL and soleus, respectively. In humans, muscles represent approximately 25% of body weight. Therefore, even a moderate increase in free fatty acid oxidation can have quantitatively important consequences on overall energy utilization.

20 In summary, these experiments show that gAcrp30 acts on muscle cells *in vivo* and *ex vivo*. The invention is drawn, *inter alia*, to the effects of gAcrp30 on muscle resulting in increased lean muscle mass and exercise capability. The increase in oleate oxidation seen in these whole muscle experiments, suggests that experiments done using the simpler C2C12 cell line model are predictive for effects *in vivo*.

25 The hindlimb muscle and liver triglyceride content was measured after gAcrp30 treatment of mice. Hind limb muscles as well as liver samples were removed from treated and untreated animals and the triglyceride and free fatty acid concentration was determined following a standard lipid extraction method (Shimabukuro et al (1997) Proc Natl Acad Sci USA 94:4637-4641; hereby incorporated herein by reference in its entirety including any figures, drawings, or tables) followed
30 by TG and FFA analysis using standard test kits.

Short-term treatment of animals with gAcrp30 (2 injections of 25 μg each given within 3 hours before sacrifice) did not change the triglyceride content either of hind limb muscle or liver tissue. However, after 3 days of treatment, during which period normal C57BL/6J mice consumed a regular rodent diet, mice that had received 25 μg of gAcrp30 twice daily showed significantly higher
35 ($p=0.002$) muscle triglyceride content (Fig.5A) than those receiving saline (control: light gray; gAcrp30: dark gray). This contrasted with a lack of increase in liver triglycerides (Fig. 5B).

Furthermore, no detectable increase in muscle TG was observed after the 16-day treatment shown independently by directly measuring the muscle TG content and by oil red O staining of frozen microscope sections. In summary, the data indicate that the increase in TG content was transient.

Ketone bodies (KB) are produced in the liver as a result of free fatty acid oxidation, but KB formation does not occur significantly in muscle. In mice receiving the high fat test meal and saline injection, the level of plasma KB increased significantly over the next 3 hours ($183 \pm 12\%$, $n=6$). Animals treated with gAcrp30, on the other hand, showed no increase in plasma KB concentrations. Although not wishing to be limited by any particular theory, this suggests that gAcrp30 inhibits either directly KB formation or can decrease KB production by inhibiting liver FFA oxidation.

Example 5: Methods of Testing the Effect of gOBG3 on Apoptosis

Prior experiments indicated that gAcrp30 either increased cell division or decreased apoptosis in C2C12 cells undergoing differentiation (Example 1). In order to assess the effect C2C12 cells can be transferred into medium containing 0.5% horse serum (rather than 2% horse serum for differentiation), which rapidly induces the onset of apoptosis (Wang & Walsh (1996) hereby incorporated herein by reference in its entirety including any figures, tables, or drawings). Cells would then be incubated for 16 hours either in the presence or absence of gACRP30 and then the level of apoptosis compared by staining for ApopTag as described in Wang & Walsh (1996).

Example 6: Methods of Assessing the Acceleration of Myoblast Differentiation with gOBG3

Prior experiments have also indicated that gAcrp30 accelerates re-organization/reorientation and differentiation of C2C12 muscle cells *in vitro* (Example 1). Rapid, highly sensitive methods are available to facilitate screening of the activity of variants of gAcrp30 (and antagonists and agonists) for their effect on differentiation. These assays include, among others, a TaqMan PCR-based method that assesses mRNA levels of muscle-specific markers of differentiation, including, but not limited to, muscle regulatory factor, myogen, alpha-actin; thermoregulatory uncoupling protein (UCP2), glucose transporter isotype glut4, myf5, beta-actin, UCP1, UCP3, and glut1 (Shimokawa et al (1998) Biochem Biophys Res Commun 246:287-292, hereby incorporated by reference herein in its entirety including any figures, drawings, or tables). Levels of mRNA would be compared between the gAcrp30 treated and the untreated cultures of differentiating C2C12 cells.

Example 7: Effect of gOBG3 on Muscle Cells *in vivo*

Experiments can be performed on normal mice as well as mice fed a high fat diet, to assess increases in the differentiation of muscle cells over a time course of treatment using the TaqMan PCR-based method described in Example 6, for example. Other muscle cell parameters discussed

herein can also be assessed over time. The animal experiments can be performed as follows, for example:

Experiment 1: 10-week-old male C57BL/6J mice were put on a very high fat/sucrose purified diet for 19 days to promote weight gain (see Example 4); the average body weight at this time was 30g. The mice were then surgically implanted with an osmotic pump (Alzet, Newark, DE) delivering either 2.5 µg/day of gAcrp30, 5 µg/day of Acrp30, or physiological saline. The mice were continued on the high fat diet and their body weight was recorded over the following 10-day period. Muscle parameters can be assessed over this same time period.

Experiment 2: mature 9 month old, male obese C57BL/6J mice that had been on the same high fat/sucrose diet for 6 months; the average body weight when the study began was 52.5±0.8g. Three groups of 8 mice were treated with saline, Acrp30 or gAcrp30 for 16 days. Animals in the treated group received twice daily 25 µg of protein subcutaneously. Body weights were recorded at appropriate time points (daily or weekly). Muscle parameters can also be assessed over the same time period.

Example 8: Mouse Models of Muscular Dystrophy

In order to further assess the efficacy of gAcrp30 and analogs, variants, and fragments thereof (as well as antagonists or agonists as needed) mouse models of muscular dystrophy can be used including, but not limited to, 129P1/ReJ-*Lama2*^{dy}, C57BL/10ScSn-*Dmd*^{mdx}/J, C57BL/6J-*Lama2*^{dy}, C57BL/6J-*Lama2*^{dy-2J}, C57BL/6Ros-*Dmd*^{mdx-2Cv}, C57BL/6Ros-*Dmd*^{mdx-3Cv}, C57BL/6Ros-*Dmd*^{mdx-4Cv}, C57BL/6Ros-*Dmd*^{mdx52Cv}, and D.B/20Ei-*Lama2*^{dy-6J}+, all of which are available from Jackson Laboratories. The experiments would include beginning to treat mice as young as possible (and then starting at increasingly later time points) with gAcrp30 as described in Example 7 and comparing the onset and severity of disease symptoms with untreated mice.

Example 9: Cellular Binding and Uptake of gOBG-3 as Detected by Fluorescence Microscopy

Fluorecein isothiocyanate (FITC) conjugation of gOBG3: Purified gOBG3 at 1 mg/mL concentration was labeled with FITC using Sigma's FluoroTag FITC conjugation kit (Stock No. FITC-1). Protocol outlined in the Sigma Handbook for small scale conjugation was followed for gOBG3 labeling.

Cell Culture: C2C12 mouse skeletal muscle cells (ATCC, Manassas, VA CRL-1772) and Hepa-1-6 mouse hepatocytes (ATCC, Manassas, VA CRL-1830) were seeded into 6 well plates at a cell density of 2x10⁵ cells per well. C2C12 and Hepa-1-6 cells were cultured according to repository's instructions for 24-48 hours prior to analysis. Assay was performed when cells were 80% confluent.

FITC labeled gOBG3 cellular binding and uptake using microscopy in C2C12 and Hepa 1-6 cells. FITC- gOBG3 (50 nM/mL) was added to each cell culture well. Cells were incubated for 1

hour at 37 °C, 5% CO₂. Cells were washed 2x with PBS, cells were scraped from well into 1 mL of PBS. Cell suspension was transferred to an eppendorf tube and centrifuged at 1000 rpm for 2 minutes. Supernatant was removed and cells resuspended in 200 µL of PBS. Binding and uptake of FITC- gOBG3 was analyzed by fluorescence microscopy under 40X magnification.

5 Analysis of C2C12 and Hepa 1-6 cells reveals identical phenotypes with respect to FITC- gOBG3 binding and uptake profiles. FITC- gOBG3 appears to be localized within vesicles in the cytoplasm of both mouse hepatocytes and mouse myoblasts, suggesting that binding and uptake of FITC-gOBG3 is occurring. Thus, this assay may be useful for identifying agents that facilitate or prevent the uptake and/or binding of OBG3 or gOBG3 polypeptides to cells by adding the substance
10 to be tested with, before, or after the FITC-gOBG3.

EXAMPLE 10: Detection of Apm-1 Fragment in Human Plasma after Immunoprecipitation

 The recombinant form of Acrp30 protein used has an apparent molecular weight of 37 kDa and forms a dimer of 74 kDa (Fig 6 A, Lane II). A proteolytic fragment that contains the entire
15 globular head region (gAcrp30) and that migrates with an apparent molecular weight of 18 kDa was generated using acetylated trypsin (Fig. 6A, lane I). Both protein preparations (Acrp30 and gAcrp30) were essentially endotoxin free; "ActiClean Etox" affinity columns were used to remove potential endotoxin contaminations (Sterogene Bioseparations Inc., Carlsbad, CA) following the manufacturer's protocol. Endotoxin levels were determined by Endosafe, Charleston, SC. As
20 determined by N-terminal sequencing of purified gAcrp30, the site of cleavage was just before amino acid 104.

 Immunoprecipitation of human plasma Apm1 followed by Western blotting was used to detect a cleavage product of Apm-1, the human homolog of Acrp30, using a globular head specific anti-serum for the immunoprecipitation step as well as for the detection step. Preimmune serum or
25 serum raised against the globular head domain or human non-homologous region (HDQETTTQGPGVLLPLPKGA) were cross-linked to protein A (Sigma Chemical CO, Saint Louis, MO) using dimethyl-pimelimidate-dihydrochloride (Sigma Chemical Co, Saint Louis, MO). After washing (0.2 M salt), proteins were eluted from protein A, were separated by SDS-PAGE, and were
30 transferred to Protran® pure nitrocellulose membrane (Schleicher and Schuell, Keene, NH) using standard procedures. Apm-1 products were visualized using globular head domain antibodies labeled with biotin; horseradish peroxidase conjugated to Streptavidin and CN/DAB substrate kit (Pierce, Rockford, IL) according to manufacturer's instructions.

 The apparent molecular weight of this truncated form was 27 kDa, corresponding to about 70% of the complete form of Apm-1 (Fig. 6B, Lane IV). This truncated form was not detectable
35 when immunoprecipitation was performed using a different antibody directed against the human non-homologous region (HDQETTTQGPGVLLPLPKGA) of Apm-1; this domain is located toward

the NH₂ terminal end of the protein outside of the globular domain (Fig. 6, Lane V). Both anti-Apm-1 antibodies directed against either the globular or the non-globular domain identified the full-length form of the protein, as well as a low abundance dimer of apparent MW 74 kDa.

5 EXAMPLE 11: Tissue Distribution of gAcrp30

Intra venous Injection of ¹²⁵I (g)Acrp30

The disappearance from plasma and subsequent tissue distribution of Acrp30 and the globular head (gAcrp30) was investigated by injecting ¹²⁵I (g)Acrp30 into CD-1 mice.

CD-1 mice were kept on regular diet with free access to food and water. The day/night light cycle was kept at 12hr ON/12hr OFF. Acrp30 and gACRP30 were prepared as previously described and labeled with ¹²⁵I to a specific activity of approximately 100 cpm/ng using Iodobeads (Pierce). The labeled protein was prepared immediately before injection and 0.1-0.5 µCi were injected through the tail vein. Clearance of the labeled protein from plasma was followed over 3 hrs after which the animals were perfused systemically with PBS + EDTA to remove all blood. Tissues were isolated (liver, adipose, kidney, skeletal muscle, brain), weighed, and the specific activity in plasma and tissue was determined.

Acrp30 showed a plasma half-life of about 7 hours. The plasma turnover of gAcrp30 was significantly faster showing a half-life close to 1 hour. The majority of the labeled Acrp30 protein was found in kidney and in skeletal muscle tissue, indicating that it represents a functional target tissue. A much smaller part of the injected protein was found in adipose and liver tissue and only a very small pool was seen in brain.

Immunohistochemistry Studies

Frozen sections of skeletal muscle from mice were stained using an antibody to the globular head of Acrp30 coupled with colorimetric detection. Briefly, C57 mice were sacrificed and various skeletal muscles were isolated. Muscles were frozen in an isopentane/liquid nitrogen bath. Tissue was then sectioned at 8 microns using a Leica 3050 cryostat and sections were picked up on microscope slides. Slides were dried overnight and fixed with 4 °C acetone after which sections were pretreated for endogenous peroxidase and biotin prior to IHC. An antibody to gAcrp30 was used on the slides at a concentration of 1:20,000. The primary antibody was detected using a biotin-labeled streptavidin system. The chromagen was DAB with hematoxylin counterstain.

Cross-sectional views of skeletal muscle show distinct outlining of muscle fibers with Acrp30 staining. A positive signal is also seen in any blood vessels in the tissue. This is not surprising since the animals were not perfused before tissues were isolated and Acrp30 should thus

be present in the plasma that would be in these vessels. This staining appears to be specific given that tissues stained with preimmune at the same concentration were clean. Also it should be noted that tissues without much Acrp30, such as liver, do not show much Acrp30 staining.

The substantial presence of Acrp30 in skeletal muscle supports the inventor's belief that
5 muscle is in fact a target tissue of this molecule.

REFERENCES

Dehouck et al. J Neurochem 54:1798-801, 1990

Shapiro, and Scherer, Curr. Biol. 8:335-338, 1998

Uysal, et al. Nature 389:610-614, 1997

5

CLAIMS

1. A method of accelerating muscle cell differentiation, comprising contacting muscle cells with gOBG3 under conditions wherein said gOBG3 binds to said cells thereby accelerating differentiation of said cells.
5
2. The method of claim 1, wherein said cells are in an individual.
3. The method of claim 2, wherein said gOBG3 is present in a pharmaceutical composition.
10
4. The method of claim 3, wherein said gOBG3 is gApm1.
5. A method of accelerating muscle cell reorganization, comprising contacting muscle cells with gOBG3 under conditions wherein said gOBG3 binds to said cells thereby accelerating reorganization of said cells.
15
6. The method of claim 5, wherein said cells are in an individual.
7. The method of claim 6, wherein said gOBG3 is present in a pharmaceutical composition.
20
8. The method of claim 7, wherein said gOBG3 is gApm1.
9. A method of increasing muscle mass in an individual, comprising contacting muscle cells in said individual with gOBG3 under conditions wherein said gOBG3 binds to said cells thereby accelerating the reorganization and differentiation of said cells and increasing said muscle mass.
25
10. The method of claim 9, wherein said gOBG3 is present in a pharmaceutical composition.
- 30 11. The method of claim 10, wherein said gOBG3 is gApm1.
12. A method of treating muscle cell disorders in an individual, comprising contacting muscle cells in said individual with gOBG3 under conditions wherein said gOBG3 binds to said cells thereby accelerating the reorganization and differentiation of said cells, and thereby treating said muscle cell disorders.
35

13. The method of claim 12, wherein said muscle cell disorders are selected from the group consisting of muscle-related eye disorders, muscle-related recovery after injuries, muscle-related recovery after surgery, muscle-related disorders of aging, muscle atrophy, muscle wasting, and muscular dystrophy.
- 5
14. The method of claim 13, wherein said gOBG3 is present in a pharmaceutical composition.
15. The method of claim 14, wherein said gOBG3 is gApm1.

1/6

	10	20	30	
1	M L L L G A V L L L L A L P G H D Q E - - - T T T Q G P G V			apmlprotein
1	M L L L Q A L L F L L I L P S H A E D D V T T T E E L A P A			adipoQprotein
1	M L L L Q A L L F L L I L P S H A E D D V T T T E E L A P A			acrp30protein2
	40	50	60	
28	L L P L P K G A C T G W M A G I P G H P G H N G A P G R D G			apmlprotein
31	L V P P P K G T C A G W M A G I P G H S G H N G T P G R D G			adipoQprotein
31	L V P P P K G T C A G W M A G I P G H P G H N G T P G R D G			acrp30protein2
	70	80	90	
58	R D G T P G E K G E K G D P G L L G P K G D I G E T G V P G			apmlprotein
61	R D G T P G E K G E K G D S G L L G P K G E T G D V G M T G			adipoQprotein
61	R D G T P G E K G E K G D A G L L G P K G E T G D V G M T G			acrp30protein2
	100	110	120	
88	A E G P R G F P G I Q G R K G E P G E G A Y V Y R S A F S V			apmlprotein
91	A E G P R G F P G T P G R K G E P G E A A Y V Y R S G F S V			adipoQprotein
91	A E G P R G F P G T P G R K G E P G E A A Y M Y R S A F S V			acrp30protein2
	130	140	150	
118	G L E T Y V T I P N M P I R F T K I F Y N Q Q N H Y D G S T			apmlprotein
121	G L E T R V T V P N V P I R F T K I F Y N Q Q N H Y D N S T			adipoQprotein
121	G L E T R V T V P N V P I R F T K I F Y N Q Q N H Y D G S T			acrp30protein2
	160	170	180	
148	G K F H C N I P G L Y Y F A Y H I T V Y M K D V K V S L F K			apmlprotein
151	G K F Y C N I P G L Y Y F S Y H I T V Y M K D V K V S L F K			adipoQprotein
151	G K F Y C N I P G L Y Y F S Y H I T V Y M K D V K V S L F K			acrp30protein2
	190	200	210	
178	K D K A M L F T Y D Q Y Q E N N V D Q A S G S V L L H L E V			apmlprotein
181	K D K A V L F T Y D Q Y Q E K N V D Q A S G S V L L H L E V			adipoQprotein
181	K D K A V L F T Y D Q Y Q E K N V D Q A S G S V L L H L E V			acrp30protein2
	220	230	240	
208	G D Q V W L Q V Y G E G E R N G L Y A D N D N D S T F T G F			apmlprotein
211	G D Q V W L Q V Y G D G D H N G L Y A D N V N D S T F T G F			adipoQprotein
211	G D Q V W L Q V Y G D G D H N G L Y A D N V N D S T F T G F			acrp30protein2
	238			
238	L L Y H D T N .			apmlprotein
241	L L F H D T - N			adipoQprotein
241	L L Y H D T - N			acrp30protein2

FIG. 1

2/6

Schematic drawing of Apm1 protein structure
(mouse homologue = Acrp30)




FIG. 2

3/6

**Oleate Oxidation in Differentiated C2C12
Cells \pm gACRP30**




Figure 3

4/6




FIG. 4

5/6




FIG. 5A

FIG. 5B

6/6




FIG. 6A

FIG. 6B

Sequence Listing

<110> GENSET

<120> Use of ACRP30 Globular Head to Promote Increases in Muscle Mass and Muscle Differentiation

<130> 87.WO1

<140>

<141>

<150> 60/239,735

<151> 2000/10/11

<150> 60/208,251

<151> 2000/5/31

<160> 7

<170> Patent.pm

<210> 1

<211> 1152

<212> DNA

<213> mus musculus

<400> 1

```

gaattcggca cgaggg atg cta ctg ttg caa gct ctc ctg ttc ctc tta atc      52
      Met Leu Leu Leu Gln Ala Leu Leu Phe Leu Leu Ile
      1          5          10
ctg ccc agt cat gcc gaa gat gac gtt act aca act gaa gag cta gct      100
Leu Pro Ser His Ala Glu Asp Asp Val Thr Thr Thr Glu Glu Leu Ala
      15          20          25
cct gct ttg gtc cct cca ccc aag gga act tgt gca ggt tgg atg gca      148
Pro Ala Leu Val Pro Pro Pro Lys Gly Thr Cys Ala Gly Trp Met Ala
      30          35          40
ggc atc cca gga cat tct ggc cac aat ggc aca cca ggc cgt gat ggc      196
Gly Ile Pro Gly His Ser Gly His Asn Gly Thr Pro Gly Arg Asp Gly
      45          50          55          60
aga gat ggc act cct gga gag aag gga gag aaa gga gat tca ggt ctt      244
Arg Asp Gly Thr Pro Gly Glu Lys Gly Glu Lys Gly Asp Ser Gly Leu
      65          70          75
ctt ggt cct aag ggt gag aca gga gat gtt gga atg aca gga gct gaa      292
Leu Gly Pro Lys Gly Glu Thr Gly Asp Val Gly Met Thr Gly Ala Glu
      80          85          90
ggg cct cgg ggc ttc ccc gga acc cct ggc agg aaa gga gag cct gga      340
Gly Pro Arg Gly Phe Pro Gly Thr Pro Gly Arg Lys Gly Glu Pro Gly
      95          100          105
gaa gcc gct tat gtg tat cgc tca ggc ttc agt gtg ggg ctg gag acc      388
Glu Ala Ala Tyr Val Tyr Arg Ser Gly Phe Ser Val Gly Leu Glu Thr
      110          115          120
cgc gtc act gtt ccc aat gta ccc att cgc ttt act aag atc ttc tac      436
Arg Val Thr Val Pro Asn Val Pro Ile Arg Phe Thr Lys Ile Phe Tyr
      125          130          135          140
aac caa cag aat cat tat gac aac agc act ggc aag ttc tac tgc aac      484
Asn Gln Gln Asn His Tyr Asp Asn Ser Thr Gly Lys Phe Tyr Cys Asn
      145          150          155
att ccg gga ctc tac tac ttc tct tac cac atc acg gtg tac atg aaa      532
Ile Pro Gly Leu Tyr Tyr Phe Ser Tyr His Ile Thr Val Tyr Met Lys
      160          165          170
gat gtg aag gtg agc ctc ttc aag aag gac aag gcc gtt ctc ttc acc      580

```


Asp	Val	Lys	Val	Ser	Leu	Phe	Lys	Lys	Asp	Lys	Ala	Val	Leu	Phe	Thr		
		175					180					185					
tac	gac	cag	tat	cag	gaa	aag	aat	gtg	gac	cag	gcc	tct	ggc	tct	gtg		628
Tyr	Asp	Gln	Tyr	Gln	Glu	Lys	Asn	Val	Asp	Gln	Ala	Ser	Gly	Ser	Val		
		190				195					200						
ctc	ctc	cat	ctg	gag	gtg	gga	gac	caa	gtc	tgg	ctc	cag	gtg	tat	ggg		676
Leu	Leu	His	Leu	Glu	Val	Gly	Asp	Gln	Val	Trp	Leu	Gln	Val	Tyr	Gly		
		205			210				215						220		
gat	ggg	gac	cac	aat	gga	ctc	tat	gca	gat	aac	gtc	aac	gac	tct	aca		724
Asp	Gly	Asp	His	Asn	Gly	Leu	Tyr	Ala	Asp	Asn	Val	Asn	Asp	Ser	Thr		
			225						230					235			
ttt	act	ggc	ttt	ctt	ctc	ttc	cat	gat	acc	aac	tga	ctg	caactac				770
Phe	Thr	Gly	Phe	Leu	Leu	Phe	His	Asp	Thr	Asn	*						
		240						245									
tcatagccca	tacaccagga	gaatcatgga	acgtcgacac	actttcagct	tagtttgaga												830
gattgatttt	attgcttagt	ttgagagtcc	tgagtattat	ccacacgtgt	actcacttgt												890
tcattaaacg	actttataaa	aaataatttg	tggttcctagt	ccagaaaaaa	aggcactccc												950
tggtctccac	gactcttaca	tggtagcaat	aacagaatga	aaatcacatt	tggtatgggg												1010
gcttcacaat	attcgcataa	ctgtctggaa	gtagaccatg	ctatttttct	gctcactgta												1070
cacaaatatt	gttcacataa	accctataat	gtaaataatga	aatacagtga	ttatcttctc												1130
aaaaaaaaact	cgtagccgaat	tc															1152

<210> 2

<211> 247

<212> PRT

<213> mus musculus

<400> 2

Met	Leu	Leu	Leu	Gln	Ala	Leu	Leu	Phe	Leu	Leu	Ile	Leu	Pro	Ser	His		
1				5				10					15				
Ala	Glu	Asp	Asp	Val	Thr	Thr	Thr	Glu	Glu	Leu	Ala	Pro	Ala	Leu	Val		
			20					25				30					
Pro	Pro	Pro	Lys	Gly	Thr	Cys	Ala	Gly	Trp	Met	Ala	Gly	Ile	Pro	Gly		
			35				40					45					
His	Ser	Gly	His	Asn	Gly	Thr	Pro	Gly	Arg	Asp	Gly	Arg	Asp	Gly	Thr		
		50			55				60								
Pro	Gly	Glu	Lys	Gly	Glu	Lys	Gly	Asp	Ser	Gly	Leu	Leu	Gly	Pro	Lys		
				70					75						80		
Gly	Glu	Thr	Gly	Asp	Val	Gly	Met	Thr	Gly	Ala	Glu	Gly	Pro	Arg	Gly		
			85					90					95				
Phe	Pro	Gly	Thr	Pro	Gly	Arg	Lys	Gly	Glu	Pro	Gly	Glu	Ala	Ala	Tyr		
			100				105						110				
Val	Tyr	Arg	Ser	Gly	Phe	Ser	Val	Gly	Leu	Glu	Thr	Arg	Val	Thr	Val		
			115				120					125					
Pro	Asn	Val	Pro	Ile	Arg	Phe	Thr	Lys	Ile	Phe	Tyr	Asn	Gln	Gln	Asn		
			130			135				140							
His	Tyr	Asp	Asn	Ser	Thr	Gly	Lys	Phe	Tyr	Cys	Asn	Ile	Pro	Gly	Leu		
				150					155						160		
Tyr	Tyr	Phe	Ser	Tyr	His	Ile	Thr	Val	Tyr	Met	Lys	Asp	Val	Lys	Val		
			165					170					175				
Ser	Leu	Phe	Lys	Lys	Asp	Lys	Ala	Val	Leu	Phe	Thr	Tyr	Asp	Gln	Tyr		
			180				185					190					
Gln	Glu	Lys	Asn	Val	Asp	Gln	Ala	Ser	Gly	Ser	Val	Leu	Leu	His	Leu		
		195					200					205					
Glu	Val	Gly	Asp	Gln	Val	Trp	Leu	Gln	Val	Tyr	Gly	Asp	Gly	Asp	His		
		210				215					220						
Asn	Gly	Leu	Tyr	Ala	Asp	Asn	Val	Asn	Asp	Ser	Thr	Phe	Thr	Gly	Phe		
		225			230				235						240		
Leu	Leu	Phe	His	Asp	Thr	Asn											
			245														

<210> 3

<211> 1276

<212> DNA

<213> mus musculus

<400> 3

```

ctctaaagat tgtcagtgga tctgacgaca ccaaaagggc tcagg atg cta ctg ttg      57
                                     Met Leu Leu Leu
                                     1
caa gct ctc ctg ttc ctc tta atc ctg ccc agt cat gcc gaa gat gac      105
Gln Ala Leu Leu Phe Leu Leu Ile Leu Pro Ser His Ala Glu Asp Asp
5                                     10                                     15                                     20
gtt act aca act gaa gag cta gct cct gct ttg gtc cct cca ccc aag      153
Val Thr Thr Thr Glu Glu Leu Ala Pro Ala Leu Val Pro Pro Pro Lys
                                     25                                     30                                     35
gga act tgt gca ggt tgg atg gca ggc atc cca gga cat cct ggc cac      201
Gly Thr Cys Ala Gly Trp Met Ala Gly Ile Pro Gly His Pro Gly His
                                     40                                     45                                     50
aat ggc aca cca ggc cgt gat ggc aga gat ggc act cct gga gag aag      249
Asn Gly Thr Pro Gly Arg Asp Gly Arg Asp Gly Thr Pro Gly Glu Lys
                                     55                                     60                                     65
gga gag aaa gga gat gca ggt ctt ctt ggt cct aag ggt gag aca gga      297
Gly Glu Lys Gly Asp Ala Gly Leu Leu Gly Pro Lys Gly Glu Thr Gly
                                     70                                     75                                     80
gat gtt gga atg aca gga gct gaa ggg cca cgg ggc ttc ccc gga acc      345
Asp Val Gly Met Thr Gly Ala Glu Gly Pro Arg Gly Phe Pro Gly Thr
85                                     90                                     95                                     100
cct ggc agg aaa gga gag cct gga gaa gcc gct tat atg tat cgc tca      393
Pro Gly Arg Lys Gly Glu Pro Gly Glu Ala Ala Tyr Met Tyr Arg Ser
                                     105                                     110                                     115
gcg ttc agt gtg ggg ctg gag acc cgc gtc act gtt ccc aat gta ccc      441
Ala Phe Ser Val Gly Leu Glu Thr Arg Val Thr Val Pro Asn Val Pro
                                     120                                     125                                     130
att cgc ttt act aag atc ttc tac aac caa cag aat cat tat gac ggc      489
Ile Arg Phe Thr Lys Ile Phe Tyr Asn Gln Gln Asn His Tyr Asp Gly
                                     135                                     140                                     145
agc act ggc aag ttc tac tgc aac att ccg gga ctc tac tac ttc tct      537
Ser Thr Gly Lys Phe Tyr Cys Asn Ile Pro Gly Leu Tyr Tyr Phe Ser
                                     150                                     155                                     160
tac cac atc acg gtg tac atg aaa gat gtg aag gtg agc ctc ttc aag      585
Tyr His Ile Thr Val Tyr Met Lys Asp Val Lys Val Ser Leu Phe Lys
165                                     170                                     175                                     180
aag gac aag gcc gtt ctc ttc acc tac gac cag tat cag gaa aag aat      633
Lys Asp Lys Ala Val Leu Phe Thr Tyr Asp Gln Tyr Gln Glu Lys Asn
                                     185                                     190                                     195
gtg gac cag gcc tct ggc tct gtg ctc ctc cat ctg gag gtg gga gac      681
Val Asp Gln Ala Ser Gly Ser Val Leu Leu His Leu Glu Val Gly Asp
                                     200                                     205                                     210
caa gtc tgg ctc cag gtg tat ggg gat ggg gac cac aat gga ctc tat      729
Gln Val Trp Leu Gln Val Tyr Gly Asp Gly Asp His Asn Gly Leu Tyr
215                                     220                                     225
gca gat aac gtc aac gac tct aca ttt act ggc ttt ctt ctc tac cat      777
Ala Asp Asn Val Asn Asp Ser Thr Phe Thr Gly Phe Leu Leu Tyr His
230                                     235                                     240
gat acc aac tga ctgcaactac ccatagccca tacaccagga gaatcatgga      829
Asp Thr Asn *
245
acagtcgaca cactttcagc ttagtttgag agattgattt tattgcttag tttgagagtc      889
ctgagtatta tccacacgtg tactcacttg ttcattaaac gactttataa aaaataatTT      949
gtgttcctag tccagaaaaa aaggcactcc ctggtctcca cgactcttac atggtagcaa      1009
taacagaatg aaaatcacat ttggtatggg ggcttcacaa tattcgcatg actgtctgga      1069
agttagccat gctatttttc tgctcactgt acacaaatat tgttcacata aaccctataa      1129
tgtaaatatg aaatacagtg attactcttc tcacaggetg agtgtatgaa tgtctaaaga      1189
cccataagta ttaaagtggg agggataaat tggaaaaaaa aaaaaaaaaa aagaaaaaact      1249
ttagagcaca ctggcgcccg ttactag      1276

```

<210> 4
 <211> 247
 <212> PRT
 <213> mus musculus

<400> 4
 Met Leu Leu Leu Gln Ala Leu Leu Phe Leu Leu Ile Leu Pro Ser His
 1 5 10 15
 Ala Glu Asp Asp Val Thr Thr Thr Glu Glu Leu Ala Pro Ala Leu Val
 20 25 30
 Pro Pro Pro Lys Gly Thr Cys Ala Gly Trp Met Ala Gly Ile Pro Gly
 35 40 45
 His Pro Gly His Asn Gly Thr Pro Gly Arg Asp Gly Arg Asp Gly Thr
 50 55 60
 Pro Gly Glu Lys Gly Glu Lys Gly Asp Ala Gly Leu Leu Gly Pro Lys
 65 70 75 80
 Gly Glu Thr Gly Asp Val Gly Met Thr Gly Ala Glu Gly Pro Arg Gly
 85 90 95
 Phe Pro Gly Thr Pro Gly Arg Lys Gly Glu Pro Gly Glu Ala Ala Tyr
 100 105 110
 Met Tyr Arg Ser Ala Phe Ser Val Gly Leu Glu Thr Arg Val Thr Val
 115 120 125
 Pro Asn Val Pro Ile Arg Phe Thr Lys Ile Phe Tyr Asn Gln Gln Asn
 130 135 140
 His Tyr Asp Gly Ser Thr Gly Lys Phe Tyr Cys Asn Ile Pro Gly Leu
 145 150 155 160
 Tyr Tyr Phe Ser Tyr His Ile Thr Val Tyr Met Lys Asp Val Lys Val
 165 170 175
 Ser Leu Phe Lys Lys Asp Lys Ala Val Leu Phe Thr Tyr Asp Gln Tyr
 180 185 190
 Gln Glu Lys Asn Val Asp Gln Ala Ser Gly Ser Val Leu Leu His Leu
 195 200 205
 Glu Val Gly Asp Gln Val Trp Leu Gln Val Tyr Gly Asp Gly Asp His
 210 215 220
 Asn Gly Leu Tyr Ala Asp Asn Val Asn Asp Ser Thr Phe Thr Gly Phe
 225 230 235 240
 Leu Leu Tyr His Asp Thr Asn
 245

<210> 5
 <211> 4517
 <212> DNA
 <213> Homo sapiens

<400> 5
 ctgattccat accagagggg ctcagg atg ctg ttg ctg gga gct gtt cta ctg 53
 Met Leu Leu Leu Gly Ala Val Leu Leu
 1 5
 cta tta gct ctg ccc ggg cat gac cag gaa acc acg act caa ggg ccc 101
 Leu Leu Ala Leu Pro Gly His Asp Gln Glu Thr Thr Gln Gly Pro
 10 15 20 25
 gga gtc ctg ctt ccc ctg ccc aag ggg gcc tgc aca ggt tgg atg gcg 149
 Gly Val Leu Leu Pro Leu Pro Lys Gly Ala Cys Thr Gly Trp Met Ala
 30 35 40
 ggc atc cca ggg cat ccg ggc cat aat ggg gcc cca ggc cgt gat ggc 197
 Gly Ile Pro Gly His Pro Gly His Asn Gly Ala Pro Gly Arg Asp Gly
 45 50 55
 aga gat ggc acc cct ggt gag aag ggt gag aaa gga gat cca ggt ctt 245
 Arg Asp Gly Thr Pro Gly Glu Lys Gly Glu Lys Gly Asp Pro Gly Leu
 60 65 70
 att ggt cct aag gga gac atc ggt gaa acc gga gta ccc ggg gct gaa 293
 Ile Gly Pro Lys Gly Asp Ile Gly Glu Thr Gly Val Pro Gly Ala Glu

75	80	85	
ggt ccc cga ggc ttt ccg gga atc caa ggc agg aaa gga gaa cct gga			341
Gly Pro Arg Gly Phe Pro Gly Ile Gln Gly Arg Lys Gly Glu Pro Gly			
90 95 100 105			
gaa ggt gcc tat gta tac cgc tca gca ttc agt gtg gga ttg gag act			389
Glu Gly Ala Tyr Val Tyr Arg Ser Ala Phe Ser Val Gly Leu Glu Thr			
110 115 120			
tac gtt act atc ccc aac atg ccc att cgc ttt acc aag atc ttc tac			437
Tyr Val Thr Ile Pro Asn Met Pro Ile Arg Phe Thr Lys Ile Phe Tyr			
125 130 135			
aat cag caa aac cac tat gat ggc tcc act ggt aaa ttc cac tgc aac			485
Asn Gln Gln Asn His Tyr Asp Gly Ser Thr Gly Lys Phe His Cys Asn			
140 145 150			
att cct ggg ctg tac tac ttt gcc tac cac atc aca gtc tat atg aag			533
Ile Pro Gly Leu Tyr Tyr Phe Ala Tyr His Ile Thr Val Tyr Met Lys			
155 160 165			
gat gtg aag gtc agc ctc ttc aag aag gac aag gct atg ctc ttc acc			581
Asp Val Lys Val Ser Leu Phe Lys Lys Asp Lys Ala Met Leu Phe Thr			
170 175 180 185			
tat gat cag tac cag gaa aat aat gtg gac cag gcc tcc ggc tct gtg			629
Tyr Asp Gln Tyr Gln Glu Asn Asn Val Asp Gln Ala Ser Gly Ser Val			
190 195 200			
ctc ctg cat ctg gag gtg ggc gac caa gtc tgg ctc cag gtg tat ggg			677
Leu Leu His Leu Glu Val Gly Asp Gln Val Trp Leu Gln Val Tyr Gly			
205 210 215			
gaa gga gag cgt aat gga ctc tat gct gat aat gac aat gac tcc acc			725
Glu Gly Glu Arg Asn Gly Leu Tyr Ala Asp Asn Asp Asn Asp Ser Thr			
220 225 230			
ttc aca ggc ttt ctt ctc tac cat gac acc aac tga tcaccactaa			771
Phe Thr Gly Phe Leu Leu Tyr His Asp Thr Asn *			
235 240 245			
ctcagagcct cctccaggcc aaacagcccc aaagtcaatt aaaggctttc agtacggtta			831
ggaagttgat tattatttag ttggaggcct ttagatatta ttcattcatt tactcattca			891
tttattcatt cattcatcaa gtaactttaa aaaaatcata tgctatgttc ccagtcctgg			951
ggagcttcac aaacatgacc agataactga ctagaaagaa gtagttgaca gtgctatttt			1011
gtgcccactg tctctcctga tgctcatatc aatcctataa ggcacaggga acaagcattc			1071
tcctgttttt acagattgta tcctgaggct gagagagtta agtgaatgtc taaggtcaca			1131
cagtattaag tgacagtgtc agaaatcaaa ccagagctgt tggactttgt tcactagact			1191
gtgccccttt atagagggtac atgttctctt tggagtgttg gtaggtgtct gtttcccacc			1251
tcacctgaga gccattgaat ttgccttcct catgaattaa aacctcccc aagcagagct			1311
tcctcagaga aagtgggtct atgatgaagt cctgtcttgg aaggactact actcaatggc			1371
ccctgcaacta ctctacttcc tcttacctat gtcccttctc atgctcttcc ctccaacggg			1431
gaaagccaac tccatctcta agtgctgaac tcatecctgt tectcaaggc cacctggcca			1491
ggagcttctc tgatgtgata tccacttttt ttcttttttg agatggagtc tcaactgttc			1551
accagggctg gagtacagtg acacgacctc ggctcactgc agcctccttc tctgggtcc			1611
aagcaattat tgtgctcag cctcccaggt agctgagact tcaggtgcat tccaccacac			1671
atggctaatt ttgtatttt tagtagaaat ggggtttcgt catgttggcc aggtgtgtct			1731
cgaactcctg gcctagggtga tccaccgcc tcgacctccc aaagtgtctg gattacaggc			1791
atgagccacc atgcccagtc gatattctcac tttttatttt gccatggatg agagtcctgg			1851
gtgtgaggaa caccctccac caggctagag gcaactgccc aggaaggact gtgcttccgt			1911
cacctctaaa tcccttgca atccttgata aatgcctcat gaagaccaat ctcttgaatc			1971
ccatatctac ccagaattaa ctccattcca gtctctgcat gtaatcagtt ttatccacag			2031
aaacattttc attttaggaa atccctgggt taagtatcaa tccttgttca gctggacaat			2091
atgaatcttt tccactgaag ttagggatga ctgtgatttt cagaacacgt ccagaatttt			2151
tcatacaagaa gtagcttga gcctgaaatg caaaacccat ggaggaattc tgaagccatt			2211
gtctccttga gtaccaacag ggtcagggaa gactgggcct cctgaattta ttattgttct			2271
ttaagaatta caggttgagg tagttgatgg tggtaaacad tctctcagga gacaataact			2331
ccagtgatgt ttttcaaga ttttagcaaa aacagagtaa atagcattct ctatcaatat			2391
ataaatttaa aaaactatct ttttgcttac agttttaaat tctgaacaat ttctcttata			2451
tgtgtattgc taatcattaa ggtattttt ttccacata taaagctttg tctttttgtt			2511
gttgtgtgtg tttttaagat ggagtttccc tctgttgcca ggctagagtg cagtggcatg			2571
atctcggtt actgcaacct ttgcctccca ggtttaagcg attcttctgc ctcagcctcc			2631
cgagtagctg ggaccacagg tgcctaccac catgccaggc taatttttgt atttttagta			2691

```

aagacagggt ttcaccatat tggccaggct ggtctcgaac tcctgacctt gtgatctgcc 2751
cgctccatt gtgtgttat ttgtgagaaa gatagatatg aggtttagag agggatgaag 2811
agggtgagagt aagccttgtg ttagtcagaa ctctgtgttg tgaatgtcat tcacaacaga 2871
aaacccaaaa tattatgcaa actactgtaa gcaagaaaaa taaaggaaaa atggaaacat 2931
ttattccttt gcataataga aattaccaga gttgttctgt ctttagataa ggtttgaacc 2991
aaagctcaaa acaatcaaga cccttttctg tatgtccttc tgttctgcct tccgcagtgt 3051
aggctttacc ctgaggtgct acacagtata gttctagggt ttccctcccg atatcaaaaa 3111
gactgtggcc tgcccagctc tcgtatcccc aagccacacc atctggctaa atggacatca 3171
tgttttctgg tgatgcccc agaggagaga ggaagctctc ttcccagat gcccagcaa 3231
gtgtaacctt gcatctcatt gctctggctg agttgtgtgc ctgtttctga ccaatcactg 3291
agtcaggagg atgaaatatt catattgact taattgcagc ttaagttagg ggtatgtaga 3351
ggatttttcc cttaaagcaaa attgggacac tgttatcaga aataggagag tggatgtag 3411
atgcaaaaata atacctgtcc acaacaaact cttaatgctg tgtttgagct ttcagtgtt 3471
tcccagagag acatagctgg aaaattccta ttgattttct ctaaaatttc aacaagtagc 3531
taaagtctgg ctatgctcac agtctcacat ctggtggggg tgggctcctt acagaacacg 3591
ctttcacagt taccctaaac tctctggggc aggttattc ctttgtggaa ccagaggcac 3651
agagacagtc aactgaggcc caacagaggc ctgagagaaa ctgaggtcaa gatttcagga 3711
ttaatgggtc tgtgatgctt tgaagtacaa ttgtggattt gtccaattct ctttagttct 3771
gtcagctttt gcttcataa ttttagcgt ctattattag atatatacat gtttagtatt 3831
atgtcttatt ggtgcattta ctctcttacc attatgtaat gtccttctt atctgtgata 3891
attttctgtg ttctgaagtc tactttgtct aaaaataaca tacgactca acttccttt 3951
ctttcttctt tcctttcttt cttccttctt ttctttctct ctctctcttt ccttccttcc 4011
ttcctctctt tctctctctc tctctctctc tctctttct tgacagactc tcgttctgtg 4071
gccctggctg gacttcagtg gtgtgatctt ggctcactgc tacctctacc atgagcaatt 4131
ctcctgcctc agcctcccaa gtagctggaa ctacaggctc atgccactgc gccagctaa 4191
tttttgtatt ttctgtagag acggggtttc accacattcg tcaggttggt ttcaaactcc 4251
tgactttgtg atccaccgc ctggcctcc caaagtgtg ggattacagg catgagccat 4311
cacacctggt caactttctt ttgattagtg tttttgtggt atatctttt ccatcatgtt 4371
actttaataa tatctatatt attgtattta aaatgtgtt cttacagact gcatgtagtt 4431
gggtataatt tttatccagt ctaaaaatat ctgtctttta attggtgtt agacaattta 4491
tatttaataa aatggtggaa tttaaa 4517

```

<210> 6

<211> 244

<212> PRT

<213> Homo sapiens

<400> 6

```

Met Leu Leu Leu Gly Ala Val Leu Leu Leu Leu Ala Leu Pro Gly His
1          5          10          15
Asp Gln Glu Thr Thr Thr Gln Gly Pro Gly Val Leu Leu Pro Leu Pro
20          25          30
Lys Gly Ala Cys Thr Gly Trp Met Ala Gly Ile Pro Gly His Pro Gly
35          40          45
His Asn Gly Ala Pro Gly Arg Asp Gly Arg Asp Gly Thr Pro Gly Glu
50          55          60
Lys Gly Glu Lys Gly Asp Pro Gly Leu Ile Gly Pro Lys Gly Asp Ile
65          70          75          80
Gly Glu Thr Gly Val Pro Gly Ala Glu Gly Pro Arg Gly Phe Pro Gly
85          90          95
Ile Gln Gly Arg Lys Gly Glu Pro Gly Glu Gly Ala Tyr Val Tyr Arg
100          105          110
Ser Ala Phe Ser Val Gly Leu Glu Thr Tyr Val Thr Ile Pro Asn Met
115          120          125
Pro Ile Arg Phe Thr Lys Ile Phe Tyr Asn Gln Gln Asn His Tyr Asp
130          135          140
Gly Ser Thr Gly Lys Phe His Cys Asn Ile Pro Gly Leu Tyr Tyr Phe
145          150          155          160
Ala Tyr His Ile Thr Val Tyr Met Lys Asp Val Lys Val Ser Leu Phe
165          170          175
Lys Lys Asp Lys Ala Met Leu Phe Thr Tyr Asp Gln Tyr Gln Glu Asn
180          185          190
Asn Val Asp Gln Ala Ser Gly Ser Val Leu Leu His Leu Glu Val Gly

```

	195		200		205
Asp	Gln	Val	Trp	Leu	Gln
	210		215		220
Tyr	Ala	Asp	Asn	Asp	Asn
225			230		235
His	Asp	Thr	Asn		

<210> 7

<211> 20966

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> 1..4811

<223> 5' regulatory region

<220>

<221> exon

<222> 4812..4851

<223> exon 1

<220>

<221> exon

<222> 15144..15365

<223> exon 2

<220>

<221> exon

<222> 16277..20559

<223> exon 3

<220>

<221> misc_feature

<222> 20560..20966

<223> 3' regulatory region

<220>

<221> allele

<222> 3787

<223> 9-27-261 : polymorphic base G or C

<220>

<221> allele

<222> 11118

<223> 99-14387-129 : polymorphic base A or C

<220>

<221> allele

<222> 15120

<223> 9-12-48 : polymorphic base C or T

<220>

<221> allele

<222> 15196

<223> 9-12-124 : polymorphic base G or T

<220>

<221> allele

<222> 15427

<223> 9-12-355 : polymorphic base G or T

<220>

<221> allele
<222> 15500
<223> 9-12-428 : polymorphic base A or G

<220>
<221> allele
<222> 15863
<223> 99-14405-105 : polymorphic base A or G

<220>
<221> allele
<222> 17170
<223> 9-16-189 : polymorphic base deletion of A

<220>
<221> primer_bind
<222> 3528..3545
<223> 9-27.pu

<220>
<221> primer_bind
<222> 3928..3946
<223> 9-27.rp complement

<220>
<221> primer_bind
<222> 10990..11008
<223> 99-14387.pu

<220>
<221> primer_bind
<222> 11423..11442
<223> 99-14387.rp complement

<220>
<221> primer_bind
<222> 15073..15092
<223> 9-12.pu

<220>
<221> primer_bind
<222> 15503..15520
<223> 9-12.rp complement

<220>
<221> primer_bind
<222> 15759..15776
<223> 99-14405.pu

<220>
<221> primer_bind
<222> 16191..16211
<223> 99-14405.rp complement

<220>
<221> primer_bind
<222> 16982..17001
<223> 9-16.pu

<220>
<221> primer_bind
<222> 17384..17402
<223> 9-16.rp complement

<220>
<221> misc_binding
<222> 3775..3799
<223> 9-27-261.probe

<220>
<221> misc_binding
<222> 11106..11130
<223> 99-14387-129.probe

<220>
<221> misc_binding
<222> 15108..15132
<223> 9-12-48.probe

<220>
<221> misc_binding
<222> 15184..15208
<223> 9-12-124.probe

<220>
<221> misc_binding
<222> 15415..15439
<223> 9-12-355.probe

<220>
<221> misc_binding
<222> 15488..15512
<223> 9-12-428.probe

<220>
<221> misc_binding
<222> 15851..15875
<223> 99-14405-105.probe

<220>
<221> misc_binding
<222> 17158..17182
<223> 9-16-189.probe

<220>
<221> primer_bind
<222> 3768..3786
<223> 9-27-261.mis

<220>
<221> primer_bind
<222> 3788..3806
<223> 9-27-261.mis complement

<220>
<221> primer_bind
<222> 11099..11117
<223> 99-14387-129.mis

<220>
<221> primer_bind
<222> 11119..11137
<223> 99-14387-129.mis complement

<220>
<221> primer_bind


```

<222> 15101..15119
<223> 9-12-48.mis

<220>
<221> primer_bind
<222> 15121..15139
<223> 9-12-48.mis complement

<220>
<221> primer_bind
<222> 15177..15195
<223> 9-12-124.mis

<220>
<221> primer_bind
<222> 15197..15215
<223> 9-12-124.mis complement

<220>
<221> primer_bind
<222> 15408..15426
<223> 9-12-355.mis

<220>
<221> primer_bind
<222> 15428..15446
<223> 9-12-355.mis complement

<220>
<221> primer_bind
<222> 15481..15499
<223> 9-12-428.mis

<220>
<221> primer_bind
<222> 15501..15519
<223> 9-12-428.mis complement

<220>
<221> primer_bind
<222> 15844..15862
<223> 99-14405-105.mis

<220>
<221> primer_bind
<222> 15864..15882
<223> 99-14405-105.mis complement

<220>
<221> primer_bind
<222> 17151..17169
<223> 9-16-189.mis

<220>
<221> primer_bind
<222> 17171..17189
<223> 9-16-189.mis complement

```

```

<400> 7
gctgatctgc tgcctcagcc ttcccaaagt gctgtaattt attaggcata agccactgtg      60
cctgcctagt gttgtacatt ctgtgggttt tgacaattgt atgcatctac atgtatgtac      120
catttatagt attcctgttt ttaatttttag ccattctagt aggcattgtag tgatatctca      180
tgggtgatttt aatttgcggt tccgtaatgg ttaataatgc tgaacatctt tgcattgtct      240

```

tgtttgtcat	ttgtgtttcc	tacttgggtga	aataattgtt	catgtccttt	gtcatttttc	300
taattgaatt	tttttttacc	atattagttt	gagatttctt	tatacaatct	agatccaaat	360
ctcttgtctc	aaatatgggt	tgcaaataca	ttcctcta	tcataatattg	ccttttcctc	420
ctcttaacag	gatgtttcag	agagcaaaag	tttttagttt	gttgaaatct	cacttttcat	480
ttttttcttt	agtggattgt	gcttttgttg	tcataatgtaa	gaactcttca	ctggccctag	540
atccttgtat	tgttttctta	agattgcat	agcaaatcac	catgaactta	gtgacaaaaa	600
gacagaaatt	tatttttca	tcctactgtg	ggcagactag	acgttaatta	ttttcatgta	660
tgctcattcc	tatgacatct	ttctgatata	ataattatag	ttattcttaa	gcttcaccct	720
tttttctatt	agctttgtta	ccttgggtgt	cactttttct	tttttgacat	tgtgacctat	780
gccagatcat	gtctgttagt	acttagcct	ccattcacct	ctccataatc	ccttttgtat	840
tcctggagct	tgatgcctga	aatgacacat	cctacattcc	tttgccagat	gggtaccagt	900
tagcttgtgc	acatgggaga	caaccgtgaa	aagactgaag	tggggaagaa	gggaggagct	960
gttgtgtttc	agttagcgcc	cctggcagtg	gcggtgacag	tggctcctgt	tcagtggcaa	1020
tgggtggagca	gctagcaaga	catgcagtaa	gcgcaggtc	ataggctatg	gtccaggagc	1080
agtcaccgat	tcctgggtct	taggcaatat	catctccctt	tgttcttcca	gcctttctaa	1140
aattattgta	ctctgactag	tacaattttt	tagtattggg	ggtagtccaa	ggacacaggc	1200
tttaaaaagt	atgaattcag	ggttgcctac	ctgcattgac	tgcgcttgaa	tcataatggc	1260
cttctgggtcg	gtggcaggag	gtgacagtcc	aaatcatgca	gtagcaaacc	agatacttaa	1320
attatcatct	gagatacttc	agaagtacag	ccgtagccat	accttcagaa	gagataaaga	1380
aatgttctcc	tggccaggcg	cgggtggctca	cgcctgtcat	tccagcactt	tgggaggccg	1440
aggggggtgga	tcacctgagg	tcgggagttc	gagaccagcc	tgaccaacat	ggggaaaccc	1500
tgtctctact	aaaaatacaa	aattagcggg	gcgtgggtggc	acatgcccct	aatcccagct	1560
actcgggagg	ctaaggcagg	ataatcgctt	gaacctgaga	ggcagagggt	gcggtgaact	1620
gagatcatgc	catagtactc	cagcctgggc	aacaagagtg	aaactccatc	tcaaaaaaaa	1680
aaaaaagaaa	aaaagataaa	gaaatgttct	cctttcttgc	catttctagg	ggtttgggga	1740
tggcgtacat	tgtctcaggg	cgtgctcact	ctaccatctt	gtcctaactc	ttatttttca	1800
aaatacagtg	cttatgcttg	gttacttcag	ttaagattat	ttttaaaaa	cataattaag	1860
caaaaatata	tggccatgct	taaacatatt	taagataaat	taagtattt	ggcctgtttc	1920
agtatcccaa	ctcacatgct	aacaggggct	tgacctgtag	ctacgggtacc	ctggaggaaa	1980
tgatcgcatt	tatttgggta	tttcgggtca	agtagtaata	gttctgtcct	gggaaaaaga	2040
ctagcctcaa	ggcaattctg	attgaatggt	tttcaattac	agtctttaaa	ccagtatggc	2100
acagaaactgg	ctctttccac	atgacggcct	ttgtgggtggg	tggcagattg	ccctgaggcc	2160
tcgcaaaatg	ctaggctttc	acaatgtcac	tgactgacag	ccaggcccag	cacagtcttg	2220
gtgtgattgt	ggggctaaag	ttattccacc	ttgtgcaata	gctacagcct	tctctaacca	2280
gctgcattct	tataaagtta	gaagaaaaata	cttttttttt	tttgagatgg	attctcgctc	2340
tgttggcccag	gctggagtg	aatgggtgca	tctcggctcg	ctgcaacctc	cgctcctgg	2400
gttcaaacga	ttctcctccc	tcagaccccc	gagtagctgg	gattgcagg	gcctgccacc	2460
acgcccggct	aacttttttg	tatttttagt	ggagacggg	tttcaccatc	ttcgtcaggc	2520
tggctctcaga	ctcctgacct	caagtgtct	gcccgcctca	gcctcccaaa	atgctgggat	2580
tacaggcatg	agctactgtg	cccggccaaa	gaaaatactt	tttatgccag	ccctgaaact	2640
acctgaagc	acatacatca	accttgaggc	ctcacactcc	atcaagagg	gtgaagggca	2700
tgaggaatta	gaaagcatag	ggatttttag	ttagacagat	ctggttcaaa	tcctagactt	2760
gtgccttgaa	caaattattt	accctcattg	aactctagat	tcattatttg	taaaatgaaa	2820
gacaataata	gttatctcca	aaggaaagtt	gaattgatc	attcatttat	tcattaatc	2880
aaactttatt	ttgtcctact	ttgtgccagg	ttctattcta	ggaactaagg	gatacaactt	2940
tgaataggca	aaatctctgc	tctcctgaag	tttacttttt	tttttttttt	ttgagacaga	3000
gtttcactct	tgtcacccag	gctggagcgc	aatgggtgctc	ttggctcact	gcaacctcca	3060
cctcctgggt	tcaagtgatt	ctcttgtctc	agcctcccaa	gtagctggga	ctacagggtat	3120
gtgccaccac	gcccggctat	ttctgcattt	ttagtagaga	tggggtttca	ccatgttggtc	3180
cagactggct	tcaaactcct	gatctcaggt	gatatgcctg	tcttggcctt	ccaaagtact	3240
gggattacag	gcctgagcca	ctgcacctga	cctgaagttt	atgttctatt	aaatagcaac	3300
agacagtaac	ataaaccaaa	aataaatagg	aaaacacat	aacaaaaatc	aaacagtgat	3360
ataattgaga	gttgcttcta	tttctttttg	ttgtcttctt	ggttcaatca	gcctgctaaa	3420
ctatatggaa	cctcattttc	atggggcact	tatttaagcc	gggggacctt	ggaaagtctc	3480
tcatgtctct	catctcaacg	gcctaattgtg	acttctcttg	aaatatattg	acattagcag	3540
gaagctgagg	ctttacatca	gatctttact	ttaatggtgg	acttgacttt	actggttagat	3600
tttttaggtc	tgtgtggact	gtggagatga	tatctggggg	gcaggcagac	acttgccctg	3660
cctctgtctg	agaaaaattct	gttttggatg	tcttgttgaa	gttgggtgctg	gcacccctaa	3720
cccttgcttg	ggtcgttaatt	taattcatca	gaattgtgtg	cttgcaagaa	ccggctcaga	3780
tcctgcscct	caaaaaacaaa	acatgagcgt	gccaaagaa	tccaagggtg	tgaatgttgc	3840
cacttcaagc	ctaaactttc	taggaacacc	taagtgggtg	gcagcttcca	gttctccagg	3900
ctgcttctag	gccagagctg	ggttccacaa	gagacagaat	aggcatatat	atgcttaagg	3960
aactggaaaa	acaggctctc	tctctctcac	aaacacacac	acacacatac	caaggtagct	4020

gtcaaaatgt	tatccgaaat	tttggaaacca	aaaaatcttg	aaagatggta	ttccaatata	4080
acatttttatg	taagttttct	attatatttag	attcaaat	cgattcgagg	ccacaagctt	4140
taagaattca	gggccttttt	aacttgccaa	gccccacacc	actccaggaa	cttcccaca	4200
ccccagttct	cagaattcat	gtgcaaggtc	tttctaaat	ccaggtcca	ggtcagagag	4260
tggaggatgt	gctctatttc	ttacctgatt	gcagaccct	ctgacagtgc	tcccttctga	4320
agcactcact	gtctgaacgt	acacagtctc	agacttaata	atgcacagtg	agcaagactg	4380
tgggtgtgata	attggcgctcc	ctgacttatt	agggcaaatc	tatgggaggg	ggagacctcc	4440
tggaccactg	agcaattaat	tcattttacat	taggaagttt	ctccgtcaga	tgcaggaaaa	4500
aaatcttgtt	ttcctgctgt	ggttttgact	tttccccat	cttctgttgc	tgttgttaga	4560
ggcaaaataa	gggtcaaggc	ctggaaacac	aagtgtttg	actgaagctc	cacttggctt	4620
ccgaagccca	agctgggttg	taccaggttc	cctaggtg	aggctgtggg	caactgccag	4680
ggacatgtgc	ctgccaccg	gcctctggcc	ctcactgagt	tggccaatgg	gaaatgacaa	4740
ttgtgagggtg	gggactgcct	gcccccgta	gtaccaggct	gttgaggctg	ggccatctcc	4800
tcctcacttc	cattctgact	gcagtctgtg	gttctgattc	cataccagag	ggtaagagca	4860
attctgtgaa	gttccaggct	gggtggggga	tgcatgcata	gcctctggct	gggatcacc	4920
aggctctccc	gtccgtagta	gtgtgggagt	ggatacagg	ggatactctg	gtcagagcag	4980
cactgtgtgga	ggcagatatg	cactgggctt	cttctccgt	tctccacag	ccccaaagga	5040
gaaaggggta	tttcagacat	tccttctaag	atgcatggaa	ccattctgaa	ttttaccag	5100
ttcgctctgt	agcaggatac	ctattgagaa	aaagttaggg	tcagtaaggt	ggaagggctt	5160
gtccacagat	gaagtccaat	tcgattaagg	gggataaggg	aatacattgt	ctcttagctt	5220
gaccaggtag	ggcaaaggaa	gaagcatata	tgaaggcagc	ttcagaaaag	tcaagctgag	5280
cactgacttc	agactggaat	taggaatcca	gctctgccac	tttattctac	tcagcaata	5340
tttactgagc	aaattctatg	ggctagacag	tggattgggt	tcacaagata	caatgagtgt	5400
gacatgggtg	ttgtctatgg	atttgggat	atatgtagg	atagggat	cttacaaggt	5460
aatcaaggag	ttctaataag	gccagccatg	gtggctcaca	cctgtaatcc	cagcaatttg	5520
ggagaccgag	gcgggtggat	cacctgaggt	caggagttcc	agactagcct	gaccaacatg	5580
gtgaaacccc	gcctctacca	aaaatacaaa	aattagttgg	gcgtgatggc	aggtgcctgt	5640
aatcccagct	tctcgggagg	ctgaggcagg	agaattgtct	gaacctggga	ggcagaggtt	5700
gcagtgagcc	gagattgttg	ccactgcaat	ccagcctggg	tgacagagcg	agactttgtg	5760
tcaaaaaaaa	aaaaaaaaag	aaagaaaaga	aaaagaggct	ctaatagag	aaaatgagaa	5820
aagcctggga	tgtagtggca	acttatgaaa	aatgttaatt	aaaaaaaac	attttctgac	5880
agaagaaaact	ggatctacct	ggtttttctg	aagcctaata	ctgctcgccc	cagttagtgc	5940
tgtttctgag	gcatcctgg	tgttttgagc	tgtggatgct	gaaggtaga	gtgggaggg	6000
tttttagaggt	taggtctgcc	cctcttgtgt	tagaggacat	ggatccctgg	tctggagagg	6060
ttctgggtttt	tggatcaagc	ctcacaaggg	gtggcaccaa	ctcactccta	ggaactccgc	6120
tagaaggaag	gccagctctg	cctaattcgg	ttggggagat	gggggtccct	ttatgctagc	6180
agaatatgtc	cgaaggagca	tgatgggtgc	agctttgttc	atgaaggcca	gtggtacaca	6240
gggagcccg	agcagtcctc	agcagtcctc	ctgcacactc	ttccttaagt	cttgaggagt	6300
cttttttttg	cacaatctca	gctcactgca	acctccgct	cccaggttca	agcgattctc	6360
ctgcctcagt	ctcccaagta	gctgagacta	caggcatg	ccaccacgcc	cagctaattt	6420
ttatattttt	agtagagatg	gggttcacca	tattggccag	gatggtctcg	atctcttgac	6480
ctcatattcc	acctgcctcg	gcctcccaaa	gtgctggtat	tacaggtgtg	agccactg	6540
cctggccgag	gagtcttaag	ctgagatcac	agcattgcac	tccagcctgg	gcaaaaagag	6600
caaaactcca	tctcaaaaaa	aaaaaaaaaa	tagacacaag	actggctcct	tgtctttttt	6660
ggggacagg	tcctcactca	tcaccaggc	tggatgcag	tggtgcaatc	acagctcact	6720
gcagcctcga	tttcccaggc	tcaagtgacc	ctcccatctt	agcctcctga	gtagctggga	6780
ctacaggtgt	gtgcaaccat	gcctggctaa	tttttaaaaa	ttttttgtag	agatgaggtc	6840
tcactatatt	ggctgggggg	cctcaaaactc	ctgggctcag	cagtcctccc	acctcagcct	6900
cccaaaaggc	tgggattata	tgcttgctct	ttttaagggtg	gctgtaggga	caaactttcc	6960
acctactcct	tgtcaagcca	gtggaccggg	gggtcccagac	atacggctaa	agtcaagagg	7020
tgatgtcttt	tggagagata	ctttcaatca	ggaatttcaa	tcagaaattc	aatcatgtgg	7080
agagagactt	atcctaataa	tgtgggtgtg	cgtgggatgc	tctgttttat	tagttccttg	7140
acagtagtga	tgtgtgtgag	tgtgtgtgtg	tgccgcgcga	cactcatttg	gatgggtgtg	7200
tatgtgtgtg	gggggtgtgt	gcgtacgtat	gtggatgtgt	ggatgtgtgt	tgtgggtgtg	7260
cgctgcata	ggtggagggtg	tgtgtatggg	tgccgggtatg	tgtgtgtgtt	gggcatggag	7320
atattgacag	ctctcccagg	gctgagtga	ggctttcggg	caaagctcct	gggagctagg	7380
caaagctgag	ttgattcctg	gttatgccat	ttattattgg	gttgacacgt	gtgaaactgc	7440
caatattcta	cactttgact	tttattttatt	tttattttta	ttttttttga	gacagagttt	7500
cacacttgct	acctaggctg	gagtgcagtg	gcgcagctc	agctcactgc	aacctctgcc	7560
tcattggattc	aagtgtattc	cctgcctcag	cctcccaagt	agctggaatt	acaggtgcc	7620
gctaccacgc	ctgactaatt	tttgtatttt	tggtagagac	gggatttcac	catgttgtcc	7680
aggctgggtct	gaaactcctg	acatcaggta	atccaccac	ctcagcctcc	caaagtgtctg	7740
ggattacagg	catgagccac	tgccgcccgc	ccattttgac	ttttaaaaa	gggagtttga	7800

tataattcaa	tccagtgggt	gaattagcta	gcacgttcc	ctctccaagt	ctcaggttct	7860
cctacacggt	agagtcaaaa	gcagggctat	gggaagatta	agtaaaataa	attttgaaaa	7920
tgcttatga	aaattacact	ccaaagaact	cgcgccagt	tcagtgttct	catgttcctc	7980
atctcacatg	atcacatttc	gcggattagg	aagctgagtc	tgagaagctc	cgtgtagtgc	8040
tttttcggag	gcaccgtgat	gtgatggaag	gctcactcgt	taggaagtca	gaacagagtc	8100
tctgagggat	catttcctta	atctgtcagt	ttcctcatct	ctgaagtggg	gctcatttcc	8160
ttccttcatg	gagttattgt	aaagatgaag	ataaataacg	tgtaaaatct	agcatgggaa	8220
ctggcttcta	taaggttcta	ataagtgcac	tcctactcct	tcccctcagc	cttcccattt	8280
gtaaaagcaa	ggcaggggtg	aggtgatttc	tggggctcct	tttggtctctg	acatttgagg	8340
attttgtatc	cttttttttt	tcagagtctt	gctctgtcac	ccaggttggg	gtgcagctca	8400
atgcaaattc	cgcctcccag	gctcaagcaa	ttcttatgtc	tcagcctcct	gagtacctgg	8460
gattacaggc	aggcaccacc	accccagct	aattttttgt	attttcagta	gagacggggt	8520
tttgccatat	tggccaggct	ggtcttgaa	tcctgacttc	atgtgaccda	cccactctag	8580
cctcccaaag	tgctgagatg	acaggtgtga	gctaccgtgc	ctggccaatt	ttgtgtgctt	8640
taatgccctt	ttctgctgga	agagttggca	ccaggtttgg	tgatctcttt	ccccacacg	8700
gctctgcctc	ctgccagtc	cagaggggac	ccgtgccttg	catttcacag	gattctgctg	8760
ttgcaactga	aattccagta	ggtcaaagtg	aaatttctca	tacacttta	catgaagata	8820
aatgatcaca	gtatggcctt	ttaggatcct	gagaacatca	cggctatccc	ctgggtataat	8880
tttaaaagca	gatgaatcca	tgctgtgctg	aggtttgcca	ggaaagccag	tgctgggatt	8940
acagtgggaag	tctttttatg	ctactttttt	cttgatcccc	tcaccccatg	gggtggcata	9000
ttgaaaggca	ggatgtgtga	ccacgatact	tttctcctcc	tgactatgt	ctaagagtct	9060
gttattgggt	tctgaagatc	agagtttaat	ttccgactcc	tctctgtgta	gctctgggat	9120
cttgaaagc	cacttaacct	ttctgaagtc	ccctttcctc	atctctaaaa	tgcatatact	9180
catcactaac	atttactgag	cactgacatg	tgcagacac	cattctaagc	attttacaca	9240
gactacacca	tttgatcttc	caacaaacag	aacactgaaa	cgcattacag	gtcagaacaa	9300
atgatttgtg	cctaagcacc	aagaccgtag	agcccggtct	ccctattcta	ccctatcctg	9360
tctctcaaaa	tgattgtgag	aatcgaatga	gacactaggt	gagaaaagg	ttttataaat	9420
agcattttta	aaatttttta	aagtccacaa	aatttttaat	tttaatacag	ataaaataga	9480
tccctttgtt	ttataaaaaag	taacaaaatt	tgttatacaa	caactatggt	atttattaat	9540
tttgcccttt	tgtatgctgc	caggaaagaa	acattaagaa	atcttaaat	gattatgggt	9600
aatcagaagg	tctgctggga	ctttttattg	ctctaactgt	acagctgac	atactacctc	9660
attttttttt	atgacacttc	aagggtgctg	ttagcttcat	cactcctctg	ttgccaaaag	9720
ctttgtgacc	aaaaacaatt	aagcagattc	ctgagtcact	aaatgacaca	taaccagagt	9780
tgagacttag	gaacttttag	tgccatgcta	agcccacagg	gacacaacaa	atagcatttt	9840
acaaaggcaa	agaattgtga	cacttgagat	ttagcttggt	gatccttgta	aaagttttct	9900
tttttaggcat	aattgagttt	tagatcatag	tactcactat	tacttagtaa	taattttttt	9960
ctgataga	tacagtgtaa	caggccgggc	gcagtggctc	atgctgttaa	tcccagcact	10020
ttgggaggcc	gagcgggg	gatcacttga	ggtcaggagt	ttgagaccag	cccggccaac	10080
atgggtgaaat	cccatctcta	ctaaaaatac	aaaaaattag	ccaggtgtgg	tcgtggattc	10140
ctgtgatccc	agctacttgg	gaggggtgag	caggagcact	agttgaaccc	aggaggcgga	10200
ggttgacgtg	agccaagatg	gtgccattgc	actccagcct	gggccacaaa	gcgagactcc	10260
acttcagaaa	caaaaaaaaa	aagagagaga	gagaaaagaa	ggaagggaag	aaggaaggaa	10320
ggaaagaagg	aaggaaggaa	ggaaagaagg	aaggaaggaa	ggaagggaag	aaggaaggaa	10380
ggaaagaagg	aaggaaggaa	agaaggaagg	aaggaaggaa	ggaagggtaa	caagcaaatg	10440
gtaacaatgg	caatatctaa	aaaaatatgt	atttttatat	gtttgtcgtt	ttatatatat	10500
gacccccact	ttagagatga	ggaaactgag	agattaagga	aacgatccct	gagagactct	10560
gttctgactt	ccaaatcggt	gagctttcca	tcgcatcacg	gtgcctccga	aagcatgaca	10620
cggagcttct	cagacttagc	ttcctaatac	gctaaacggg	attatgtgag	atgaggaaca	10680
tgagaacgct	gacatgggtg	agggttcctt	ggagtatcat	tttcatgtgg	cattttcaaa	10740
acttatttta	cctaattctt	ccaaagccct	gcttttgact	ctaattgtgc	tctgagact	10800
tgagagcgc	aagatgctag	cgacagagca	agactccatc	tccagataaa	taaataagta	10860
aaataaaaaa	gaacacaaat	aattttgaaa	attttttga	aaattaggca	cgtttgcact	10920
gaccttcaat	gtttattaat	tgctggtttc	ccaccagaa	ttaagttgga	atgcaacttt	10980
cttttacaat	cagagtccgt	tcttggtctt	ggaaacttct	gaggctcctg	tgctaatecc	11040
actcttgat	ttttggcacc	tctacccctg	gccactgtca	tggaaccag	gctgatcgca	11100
cctattagtg	gagaaatmtg	tccataatac	tgaagtttgg	ggacaaacag	tgttccctta	11160
gggtaggaga	aagagatctt	tatttttaac	aaagggggag	gagccagaaa	actccagaga	11220
cccctgagtg	tgccctctct	ccaaggtttg	gggtaagccc	cccgtcaccc	tttatctctg	11280
gggctttcac	atatctctga	ttctctcttc	ctgtttccca	gcagaaaagg	atggagcctc	11340
acagattctt	cctattctctg	gagaaaaaca	tgcatggagc	tcaaagttct	tctcaggagt	11400
tttattgcca	aagccataat	aagaaagggt	ggaggtgaca	agcagtgagg	aagtttaaa	11460
atgcatgaaa	tctgtaaagt	ctcagaacaa	gaattctcct	aaaatgcaa	aggggctttg	11520
ctggctcccc	cttggtctct	catgtagctc	acctcttttt	tcttatcttg	agactagtca	11580

aacctaagct	gtttctcatt	ttatttccag	aagctattga	gaacactctc	ctgaattctt	11640
caaatccagt	agagggcgac	aaatgtacat	ataaatgatg	gtagtgggtc	ttaaataaag	11700
actcatgaca	cctaaagggg	cagcacctga	gtctgattgc	acctgtttct	gttgctgttt	11760
ctgtctctct	tctctctgtc	tgccatttca	ttatcaatgg	ttactttact	tataagatca	11820
tattagaacc	tgatatttga	taaatgatgc	atcagatcta	tagtgagaga	aaaaattaat	11880
gcaattaaag	gtgttgtaac	agctagtctt	caagtgggga	gaaatcattt	gagtacctta	11940
ggtcacagct	tacatcāaaa	caaaaaatca	gagctacatt	aaaaagtga	attttaacta	12000
tatcaaacia	tagaaaaaaa	cagaagaaaa	ttgaatactt	actaaatctt	agcatgaata	12060
agaactgttt	aacacttaga	ggcaaggact	ggcggtgggtg	gctcatgctt	ttaatccag	12120
gactttggga	ggccaaggcg	ggcggtacac	ctgaggtcag	gagtttgaga	ctagcctggc	12180
caacatgggtg	aaaccccgtc	tctactaaaa	aatgcaaaaa	ttagctgcgt	gtgggtgggtg	12240
atgcctgtaa	tctcagctac	ttgggaggct	aaggcatgag	aatcgcttga	acctgggagg	12300
tggaggctgt	agtgagccga	gattgtgcc	ctgcactaca	gcctgggtga	cagtgtgaaa	12360
tcctctctct	caaaaaaaa	aaaaaaaaa	agcaactag	agcagtgagg	taccattatt	12420
tcctttgctc	actaaactga	caacacacaa	atgtttttta	taatacccaa	agctgatgag	12480
ggtagttaag	gtatgccctt	ttatacacac	actaatgatg	tactactggg	tggcagata	12540
acatatggctg	ccatgtgggg	atatgtatca	ggagacttaa	aaatgtgcat	accttttggg	12600
ccagtaattt	acttctggga	atctgtcata	acagaataat	aatcttgggg	aaagctacat	12660
gcctaaggat	attttaaata	ttatttāaaa	atcaaagtat	aatttcttac	agaatataaa	12720
ataatatttt	aaaatgaaaa	tatgctāaaa	gtttgatgaa	atataaatgg	tcaaatatat	12780
attgattata	tcactttact	agactagcac	tcactctgag	acgttāaaaa	tagtcattat	12840
aaaaactaga	aatgccaāaa	gacaaaataa	aggaataaag	ttttacataa	agtatgattc	12900
cactatgttt	aaaaataaac	agagacattc	ttggagttga	gtattgtttt	cttttctgtc	12960
atgtccaaag	aactatataa	ctattatttt	taatgaacta	tatatgtaat	atatacatat	13020
agtttatatg	tatatācaaa	atttatctca	tatatatgat	aaagatgaaa	gatgagttgg	13080
atgtgccacg	tgaagtgggt	agtatāgaaa	cccaggtaat	ggggcatagg	agtgggattc	13140
cagataccag	gcccattgtt	ttggggtgag	attgccaatc	acggctcttc	ttccatccct	13200
cacagaggag	taggtttgtc	ttcaacaaac	cttcagttgt	cctgaagaca	aacctaattc	13260
tggagacttc	atataatcta	gaagagācaa	gcaactgat	gaaaaatagt	gaatttttaa	13320
ggtāaaataa	agtacatgga	ctacactttg	tttagaatca	gattcttggg	attaaccaca	13380
ttaaccacāa	gagggcttta	gtgatgcctc	taatccagga	tcctaaggacc	tatttctctc	13440
tgtgagatgc	tttctcccaa	ctccttgggtg	agagtgggaa	gactaagacc	tcagcaatct	13500
gaggtggagg	cctaagatcc	ccetaagatc	ggaggcagaa	tctgagaggg	gataaaagtc	13560
cctataccctg	tattgggccc	ttttctggga	gggggatatc	aaagaatgat	tttgagacag	13620
ggaggctttt	gactacctgt	gccacttgag	ctctttgcta	gggctccaga	atacatattt	13680
caaatācat	ccccctccct	ccttctctcc	ctcttccact	cttctctttt	atcttctctt	13740
cttcttttcc	ttctctcttc	ccttctcttc	tctggctctc	tcatgatttc	ttttctctat	13800
tataaaagtgt	cttatttagt	ccttactctg	cttattagtg	gttagtcttt	gtcccctggg	13860
acttgctgtt	taatggagaa	atgggtgagc	aaaacagaaa	ttacagcaga	gtgcaataat	13920
agagctaagc	caggtgtata	aatccattct	cacactgctg	tāaaaaacta	ctgggttaatt	13980
tataaagaaa	agaggtttta	ttgactcaca	gttccacagg	ctgtacagga	agcatggctg	14040
gggaggcctc	agāaaactta	caatcatggg	ggaagaaaga	gcgaagggga	agcaagcaca	14100
tcacacagca	gcaggagaga	gagagagaaa	gagagagaga	gagaatatag	gggaagtgtc	14160
acacactttc	aaccagatct	tgtgagaatt	cacctactat	catgagaaca	gcaagggata	14220
agtctgcctc	catgattcag	tcacctccta	ccaggccctc	tctccaacac	atgtcgacgt	14280
gctattttggg	tggggacaca	gacccāaac	atattaccag	ggcactggag	aaacacagag	14340
gggāaaaga	cagccaagga	gtgagatgga	gaacaaggag	gacttcttga	aacagatgac	14400
atccāaaactg	ggtcctgaaa	gctgaataga	gattagacag	gggaggaggg	gcagctaaag	14460
atggctcagg	caāacāaagg	gccaggggat	atgttcatgg	gatgatgtgt	ctctcgttgt	14520
ctgcttaaca	caaggtgagt	ctctccctcc	ctctctctct	ctttttctct	gtgtgtgttt	14580
gtgtgtgtgc	atgtgtgcaa	atgtaatata	cccaatagtc	aaacatgtgc	cccaggagag	14640
gggtagagga	agaaagagaa	tgagagagta	agaaggagga	atagacacag	aaaatgagag	14700
agaagggggg	aagāaaagg	aagaaaggag	ccagaggaga	gaagctgggt	agcattgaat	14760
ggagcaatct	gtgtcatcgt	acttgggāaa	cccaaggatg	gattcttggc	aagtcgactc	14820
ttggagcttt	ccctgtgctt	ggtcctgtgc	tcagacatgg	gāaaattaga	ggagtgtcat	14880
ctgtgcaatc	actgaattca	taatcttggg	gaggāaaagg	gactacacac	agggāataat	14940
gctaagtatt	acagatttca	gggcagāaag	agatcaaggg	gggctgcaat	attcagāaaa	15000
gtcttctctg	aaaagttaga	tacttagāaa	gcagctccta	gaagtagact	ctgctgagat	15060
ggacggagtc	ctttgtaggc	cccaactggg	tgtgtgtgtg	gggtctgtct	ctccatggcy	15120
gacagtgcac	atgtggattc	cagggtcag	gatgtctgtg	ctgggagctg	ttctactgct	15180
attagctctg	cccgkcatg	accaggāaac	cacgactcaa	gggcccggag	tcctgcttcc	15240
cctgccaag	ggggcctgca	caggttggat	ggcgggcatc	ccagggcatc	cggggcataa	15300
tggggcccca	ggcctgtgatg	gcagagatgg	caccctgggt	gagaagggtg	agaaaggaga	15360

tccaggttaag	aatgtttctg	gcctctttca	tcacagacct	cctacactga	tataaactat	15420
atgaagkcat	tcattattaa	ctaaggccta	gacacagggg	gaaagcaaa	cttttttatg	15480
ttaaccataa	gcaacctgar	gtgatttggg	gttggctctc	caaggatgag	tgtagatggt	15540
gcctctataa	ccaagacttt	ggctttgtcg	catctgcagc	tccttttcca	tcccttttcc	15600
catcttcacc	ctcatcccta	ttcccagtac	attcatattc	tgattcctct	ttctgtctgc	15660
ttaaactcca	tttcacccag	tggcattcaa	ccacatttac	tgcacacccc	ctgaaaggct	15720
cagtcctgcc	tttggggaac	tcttgatcta	ggtaagatgt	ctaattgtga	aggctctggt	15780
ggtggttacc	acaagaaagt	ctactctaaa	aatgtcaaac	tgaatgtgaa	caagtattca	15840
aagtatggag	catagagaaa	atrtactcac	cgtggacctg	atgaagaatg	aaggcttcaa	15900
ggaggaggca	gagcttcagc	taggccttga	atgatgggta	ggcagaatag	aggaggagag	15960
acatcctaga	tggagggggt	agaattgcaa	aaccagggtt	gatggtgcca	gcacataaag	16020
ggctggcagg	gtggagggtc	tatgatagag	acctatagga	gataaagata	gagttgaaat	16080
tatgggagcc	tccatgtctg	tgggagatat	agaaggagga	ggtaacacct	ctctcctttt	16140
gggagctctt	attggtttct	tgatctataa	gtcaagaagg	ttgtgagtgg	gagccacagg	16200
gatggtaatt	taggctgtaa	ccaacctagg	caggagttct	gttctttgta	gtcactgagg	16260
tcttctcatt	ccttaggtct	tattggtcct	aagggagaca	tcggtgaaac	cggagtaccc	16320
ggggctgaag	gtccccagg	ctttccggga	atccaaggca	ggaaaggaga	acctggagaa	16380
ggtgcctatg	tataccgctc	agcattcagt	gtgggattgg	agacttacgt	tactatcccc	16440
aacatgccca	ttcgctttac	caagatcttc	tacaatcagc	aaaaccacta	tgatggctcc	16500
actggtaaat	tccactgcaa	cattcctggg	ctgtactact	ttgcctacca	catcacagtc	16560
tatatgaagg	atgtgaagg	cagcctcttc	aagaaggaca	aggctatgct	cttcacctat	16620
gatcagttacc	aggaaaataa	tgtggaccag	gcctccggct	ctgtgctcct	gcatctggag	16680
gtgggcgacc	aagtctggct	ccagggtgat	ggggaggagg	agcgtaatgg	actctatgct	16740
gataatgaca	atgactccac	cttcacaggc	tttcttctct	accatgacac	caactgatca	16800
ccactaaact	agagcctcct	ccaggccaaa	cagcccaaaa	gtcaattaaa	ggctttcagt	16860
acgggttagga	agttgattat	tatttagttg	gaggccttta	gatattattc	attcattttac	16920
tcattcattt	attcattcat	tcattcaagta	acttttaaaa	aatcatatgc	tatgttccca	16980
gtcctgggga	gcttcacaaa	catgaccaga	taactgacta	gaaagaagta	gttgacagtg	17040
ctattttgtg	cccactgtct	ctcctgatgc	tcatatcaat	cctataaggc	acagggaaca	17100
agcattctcc	tgtttttaca	gattgtatcc	tgaggctgag	agagtttaagt	gaatgtctaa	17160
ggtcacacaa	gtatttaagtg	acagtgtag	aaatcaaacc	cagagctgtg	gactttgttc	17220
actagactgt	gcccttttat	agaggtaacat	gttctctttg	gagtgttgg	aggtgtctgt	17280
ttcccacctc	acctgagagc	cattgaattt	gccttctctc	tgaattaaaa	cctcccccaa	17340
gcagagcttc	ctcagagaaa	gtggttctat	gatgaagtcc	tgtcttgga	ggactactac	17400
tcaatggccc	ctgcactact	ctacttccctc	ttacctatgt	cccttctcat	gcctttccct	17460
ccaaagggga	aagccaactc	catctctaag	tgtgaaactc	atccctgttc	ctcaaggcca	17520
cttggccagg	agcttctctg	atgtgatatc	cacttttttt	tttttttgag	atggagtctc	17580
actctgtcac	ccaggctgga	gtacagtgc	acgacctcgg	ctcactgcag	cctccttctc	17640
ctgggtccaa	gcaattattg	tgcctcagcc	tcccagtag	ctgagacttc	aggtgcattc	17700
caccacacat	ggctaatttt	tgtattttta	gtagaaatgg	ggtttcgtca	tgttgccag	17760
gctggtctcg	aactcctggc	ctagggtgatc	caccgcctc	gacctcccaa	agtgtctgga	17820
ttacaggcat	gagccaccat	gccagtcga	tatctcactt	tttattttgc	catggatgag	17880
agtcttgggt	gtgaggaaca	cctcccacca	ggctagaggc	aactgcccag	gaaggactgt	17940
gcttccgtca	cctctaaatc	ccttgcatga	ccttgataaa	tgctctcatga	agaccaatct	18000
cttgaatccc	gtatctaccc	agaattaaact	ccattccagt	ctctgcatgt	aatcagtttt	18060
atccacagaa	acattttcat	tttaggaaat	ccctggtttt	aagtatcaat	ccttggttcag	18120
ctggacaata	tgaatctttt	ccactgaagt	tagggatgac	tgtgattttc	agaacacgtc	18180
cagaattttt	catcaagaag	gtagcttgag	cctgaaatgc	aaaacccatg	gaggaaattct	18240
gaagccattg	tctccttgag	taccaacagg	gtcagggaag	actgggcctc	ctgaattttat	18300
tattgttctt	taagaattac	aggttgaggt	agttgatggt	ggtaaacatt	ctctcaggag	18360
acaataaact	cagtgtatgt	cttcaaagat	tttagcaaaa	acagagtaaa	tagcattctc	18420
tatcaatata	taaatattaa	aaactatctt	tttgcttaca	gttttaaatc	ctgaacaatt	18480
ctctcttaca	tgtgtattgc	taatcattaa	ggattatttt	tttccacata	taaagctttg	18540
tctttttgtt	gttgttgttg	tttttaagat	ggagtttccc	tctgttgcca	ggctagagtg	18600
cagtggcatg	atctcggctt	actgcaacct	ttgcctccca	ggttcaagcg	attcttctgc	18660
ctcagcttcc	cgagttagctg	ggaccacagg	tgccctaccac	catgccaggc	taatttttgt	18720
atttttagta	aagacagggt	ttcaccat	tggccaggct	ggtctcgaac	tcttgacctt	18780
gtgatctgcc	cacctccatt	tttgttgtta	ttttttgaga	aagatagata	tgaggtttag	18840
agagggatga	agaggtgaga	gtaagccttg	tgtagtcag	aactctgtgt	tgtgaatgtc	18900
attcacaca	gaaaacccaa	aatattatgc	aaactactgt	aagcaagaaa	aataaaggaa	18960
aatgggaaac	atttattcct	ttgcataata	gaaattacca	gagttgttct	gtcttttagat	19020
aaggtttgaa	caaagctca	aaacaatcaa	gaccttttct	tgtatgtcct	tctgttctgc	19080
cttccgcagt	gtaggcttta	ccctcaggtg	ctacacagta	tagttctagg	gtttccctcc	19140

cgatatcaaa	aagactgtgg	cctgcccagc	tctcgtatcc	ccaagccaca	ccatctggct	19200
aaatggacat	catgttttct	ggtgatgccc	aaagaggaga	gaggaagctc	tctttcccag	19260
atgccccagc	aagtgttaacc	ttgcatctca	ttgctctggc	tgagtttgt	gcctgtttct	19320
gaccaatcac	tgagtcagga	ggatgaaata	ttcatattga	cttaattgca	gcttaagtta	19380
ggggtatgta	gaggtatttt	ccctaaagca	aaattgggac	actgttatca	gaaataggag	19440
agtggatgat	agatgcacaa	taatacctgt	ccacaacaaa	ctcttaatgc	tgtgtttgag	19500
ctttcatgag	tttcccagag	agacatagct	ggaaaattcc	tattgatttt	ctctaaaatt	19560
tcaacaagta	gctaaagtct	ggctatgctc	acagtctcac	atctgggttg	ggtgggctcc	19620
ttacagaaca	cgctttcaca	gttaccctaa	actctctggg	gcagggttat	tcctttgtgg	19680
aaccagaggc	acagagagag	tcaactgagg	ccaaaagagg	cctgagagaa	actgaggcca	19740
agatttcagg	attaatggtc	ctgtgatgct	ttgaagtaca	attgtggatt	tgtccaattc	19800
tcttttagttc	tgctcagctt	tgcttcatat	attttagcgc	tctattatta	gatataata	19860
tgtttagtat	tatgtcttat	tggtgcattt	actctcttat	cattatgtaa	tgtccttctt	19920
tatctgtgat	aattttctgt	gttctgaagt	ctactttgtc	taaaaaatac	atacgcactc	19980
aacttccttt	tctttcttcc	ttcctttctt	tcttccttcc	tttctttctc	tctctctctc	20040
tttcttctct	tccttcctcc	ttttctttct	ctctctctct	ctctctcttt	ttttgacaga	20100
ctctogttct	gtggccctgg	ctggagttca	gtgggtgat	cttggctcac	tgctacctct	20160
accatgagca	attctcctgc	ctcagcctcc	caagtagctg	gaactacagg	ctcatgccac	20220
tgcgcccagc	taatttttgt	atttttcgta	gagacggggt	ttcaccacat	tcgtcagggt	20280
ggtttcaaac	tcctgacttt	gtgatccacc	cgcctcgccc	tcccaaagtg	ctgggattac	20340
aggcatgagc	catcacacct	ggtcaacttt	cttttgatta	gtgtttttgt	ggtatatctt	20400
tttccatcat	gttactttta	atatacttat	attattgtat	ttaaaatgtg	tttcttacag	20460
actgcatgta	gttgggtata	atttttatcc	agtctaaaaa	tatctgtctt	ttaattgggt	20520
tttagacaat	ttatatttta	taaaattggt	gaatttaaga	tggatgactg	ttttatttgt	20580
ttgtgtttca	ccactttctg	tttattctct	ttccagaatt	cttttggtatt	gtttaaatat	20640
ttcataatat	tttatcttaa	tttattttat	gggtatttgc	ctatatctct	ttgtgggtatt	20700
ttttagtggg	tgcttgaggg	attacaatgt	acttaacttt	tcacagtgtg	cataaagtta	20760
atattttgcc	acttgacagta	aaccgtagaa	ggcttataat	catattagta	cctctatcca	20820
ctttctttta	tgttgtagtt	gtcatatata	ttacatctat	atacactgaa	acattatagg	20880
caatgttatg	attttttgcat	tcgtcagtc	tatatatatt	ttaaagaatt	taagaggaga	20940
aaaatacata	ttcagatatt	catcat				20966

1

1